

GALERKIN FINITE ELEMENT METHOD FOR GENERALIZED FORCHHEIMER EQUATION OF SLIGHTLY COMPRESSIBLE FLUIDS IN POROUS MEDIA

THINH KIEU [†]

Abstract. We consider the generalized Forchheimer flows for slightly compressible fluids. Using Muskat's and Ward's general form of Forchheimer equations, we describe the fluid dynamics by a nonlinear degenerate parabolic equation for the density. We study Galerkin finite elements method for the initial boundary value problem. The existence and uniqueness of the approximation are proved. The prior estimates for the solutions in $L^\infty(0, T; L^q(\Omega))$, $q \geq 2$, time derivative in $L^\infty(0, T; L^2(\Omega))$ and gradient in $L^\infty(0, T; W^{1, 2-a}(\Omega))$, with $a \in (0, 1)$ are established. Error estimates for the density variable are derived in several norms for both continuous and discrete time procedures. Numerical experiments using backward Euler scheme confirm the theoretical analysis regarding convergence rates.

Key words. Porous media, immersible flow, error analysis, Galerkin finite element, nonlinear degenerate parabolic equations, generalized Forchheimer equations, numerical analysis.

AMS subject classifications. 65M12, 65M15, 65M60, 35Q35, 76S05.

1. Introduction . Fluid flow through porous media has been studied in many fields such as chemistry, physics, engineering, and geology. The most common equation to describe fluid flows in porous media is the Darcy law

$$-\nabla p = \frac{\mu}{\kappa} v, \quad (1.1)$$

where p , v , μ , κ are, respectively (resp.), the pressure, velocity, absolute viscosity and permeability.

For the Reynolds number large, for instance the high-velocity flow [3, 30], Darcy's law is not valid any more. A nonlinear relationship between velocity and gradient of pressure is introduced, as suggested by Forchheimer in [15, 16], by adding the higher order term of velocity to the Darcy law. It is known as generalized Forchheimer laws, see [3, 30, 31, 34, 36] and references therein. Forchheimer established the following three nonlinear empirical models:

$$-\nabla p = av + b|v|v, \quad -\nabla p = av + b|v|v + c|v|^2v, \quad -\nabla p = av + d|v|^{m-1}v, m \in (1, 2). \quad (1.2)$$

Above, the positive constants a, b, c, d are obtained from experiments.

The generalized Forchheimer equation of (1.1) and (1.2) were proposed in [2, 17, 18] of the form

$$-\nabla p = \sum_{i=0}^N a_i |v|^{\alpha_i} v. \quad (1.3)$$

These equations are analyzed numerically in [8, 26, 32], theoretically in [2, 18–21, 24] for single phase flows, and also in [22, 23] for two phase flows.

In order to take into account the the dependence on density in generalized Forchheimer equation, we modify (1.3) using dimension analysis by Muskat [30] and Ward [36]. They proposed the following equation for both laminar and turbulent flows in porous media:

$$-\nabla p = F(v^\alpha \kappa^{\frac{\alpha-3}{2}} \rho^{\alpha-1} \mu^{2-\alpha}), \text{ where } F \text{ is a function of one variable.} \quad (1.4)$$

In particular, when $\alpha = 1, 2$, Ward [36] established from experimental data that

$$-\nabla p = \frac{\mu}{\kappa} v + c_F \frac{\rho}{\sqrt{\kappa}} |v|v, \quad \text{where } c_F > 0. \quad (1.5)$$

[†]Department of Mathematics, University of North Georgia, Gainesville Campus, 3820 Mundy Mill Rd., Oakwood, GA 30566, U.S.A. (thinh.kieu@ung.edu).

Combining (1.3) with the suggestive form (1.4) for the dependence on ρ and v , we propose the following equation

$$-\nabla p = \sum_{i=0}^N a_i \rho^{\alpha_i} |v|^{\alpha_i} v, \quad (1.6)$$

where $N \geq 1$, $\alpha_0 = 0 < \alpha_1 < \dots < \alpha_N$ are real numbers, the coefficients $a_0, \dots, a_N$ are positive. Here, the viscosity and permeability are considered constant and we do not specify the dependence of a_i 's on them.

Multiplying both sides of previous equation to ρ , we obtain

$$g(|\rho v|) \rho v = -\rho \nabla p, \quad (1.7)$$

where the function g is a generalized polynomial with non-negative coefficients. More precisely, the function $g : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is of the form

$$g(s) = a_0 s^{\alpha_0} + a_1 s^{\alpha_1} + \dots + a_N s^{\alpha_N}, \quad s \geq 0, \quad (1.8)$$

where $N \geq 1$, $\alpha_0 = 0 < \alpha_1 < \dots < \alpha_N$ are fixed real numbers, the coefficients $a_0, \dots, a_N$ are non-negative numbers with $a_0 > 0$ and $a_N > 0$.

The equation of state which, for slightly compressible fluids, is

$$\frac{1}{\rho} \frac{d\rho}{dp} = \frac{1}{\kappa} = \text{const.} > 0.$$

Hence

$$\nabla \rho = \frac{1}{\kappa} \rho \nabla p, \quad \text{or} \quad \rho \nabla p = \kappa \nabla \rho. \quad (1.9)$$

Combining (1.8) and (1.9) implies that

$$g(|\rho v|) \rho v = -\kappa \nabla \rho. \quad (1.10)$$

By rescaling coefficients of g we have

$$g(|\rho v|) \rho v = -\nabla \rho.$$

Hence

$$\rho v = -K(|\nabla \rho|) \nabla \rho, \quad (1.11)$$

where the function $K : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is defined for $\xi \geq 0$ by

$$K(\xi) = \frac{1}{g(s(\xi))}, \quad \text{with } s = s(\xi) \text{ being the unique non-negative solution of } sg(s) = \xi. \quad (1.12)$$

The continuity equation is

$$\phi \rho_t + \text{div}(\rho v) = f. \quad (1.13)$$

where constant $\phi \in (0, 1)$ is the porosity, f is external mass flow rate.

By combining (1.11) and (1.13) we obtain

$$\phi \rho_t - \nabla \cdot (K(|\nabla \rho|) \nabla \rho) = f. \quad (1.14)$$

Then by scaling the time variable in (1.14) implies

$$\rho_t - \nabla \cdot (K(|\nabla \rho|) \nabla \rho) = f. \quad (1.15)$$

The system of equations describing the fluid motion reduces to a scalar equation of density function. This is a nonlinear parabolic equation which degenerates as the gradient of density approach to infinity.

For the existence and regularity theory of degenerate parabolic equations, see e.g. [7, 19, 27]. The popular numerical method for modeling flow in porous media are mixed finite element method. Arbogast, Wheeler and Zhang [1] first analyzed mixed finite element approximations of degenerate parabolic equation arising in flow in porous media. Not so long later, Woodward and Dawson in [37] study of expanded mixed finite element methods for a nonlinear parabolic equation modeling flow into variably saturated porous media. In recently years, the Galerkin finite element method for a coupled nonlinear degenerate system of advection-diffusion equations has studied in [9–14]. In their analysis, the Kirchhoff transformation is used to move the nonlinearity from coefficient K to the gradient and thus simplifies analysis of the equations. This transformation does not applicable for the equation (1.15).

In this paper, we combine techniques in [17–21, 24] utilizing the special structures of equation to obtain the error estimates for the solutions in several norms of interest. We obtain these results without any extra regularity assumption, though the order of error estimates are far from optimal order.

The paper is organized as follows: In §2 we introduce notations, relevant results in [2, 17, 26] and suitable trace estimates in Lemma 2.4 for the nature of our equation. In §3 we consider the semidiscrete finite element Galerkin approximation and an implicit backward difference time discretization to solve problem (3.1). In §4 we establish many bounds for solutions, its time derivative and gradient to problem (3.2) and (3.4) in Lebesgue norms. In §5 we analyze two version of a Galerkin finite element approximations, the continuous Galerkin method and the discrete Galerkin method. Using the monotonicity properties of Forchheimer equation and the boundedness of solutions, the *priori* error estimates are derived for solution in L^q -norm with $2 \leq q \leq \infty$ and for gradient of solution in L^{2-a} -norm. Finally, in §6 numerical examples using the Lagrange elements of order 2 are carried out in two-dimension. The results strongly support our theoretical analysis regarding convergence rates.

2. Notations and preliminary results. Suppose that Ω is an open, bounded subset of $\mathbb{R}^d$, with $d = 2, 3, \dots$, and has C^1 -boundary $\Gamma = \partial\Omega$. Let $L^2(\Omega)$ be the set of square integrable functions on Ω and $(L^2(\Omega))^d$ the space of d -dimensional vectors which have all components in $L^2(\Omega)$. We denote $(\cdot, \cdot)$ the inner product in either $L^2(\Omega)$ or $(L^2(\Omega))^d$ that is

$$(\xi, \eta) = \int_{\Omega} \xi \eta dx \quad \text{or} \quad (\boldsymbol{\xi}, \boldsymbol{\eta}) = \int_{\Omega} \boldsymbol{\xi} \cdot \boldsymbol{\eta} dx.$$

The notation $\langle \cdot, \cdot \rangle$ will be used for the $L^2(\partial\Omega)$ inner-product and $\|u\|_{L^p} = \|u\|_{L^p(\Omega)}$ for standard Lebesgue norm of the measurable function. The notation $\|\cdot\|$ will means scalar norm $\|\cdot\|_{L^2(\Omega)}$ or vector norm $\|\cdot\|_{(L^2(\Omega))^d}$. We denote $\|u\|_{L^p(L^q)} = \|u\|_{L^p(0,T;L^q(\Omega))}$, $1 \leq p, q < \infty$ means the mixed Lebesgue norm for a function u while $\|u\|_{L^p(H^q)} = \|u\|_{L^p(0,T;H^q(\Omega))}$, $1 \leq p, q < \infty$ stand for the mixed Sobolev-Lebesgue norm of a function u .

For $1 \leq q \leq +\infty$ and m any nonnegative integer, let

$$W^{m,q}(\Omega) = \{u \in L^q(\Omega), D^q u \in L^q(\Omega), |q| \leq m\}$$

denote a Sobolev space endowed with the norm

$$\|u\|_{m,q} = \left(\sum_{|\alpha| \leq m} \|D^\alpha u\|_{L^q(\Omega)}^q \right)^{\frac{1}{q}}.$$

Define $H^m(\Omega) = W^{m,2}(\Omega)$ with the norm $\|\cdot\|_m = \|\cdot\|_{m,2}$.

Throughout this paper, we use short hand notations,

$$I = (0, T), \quad \|\rho(t)\| = \|\rho(\cdot, t)\|_{L^2(\Omega)}, \forall t \in I \quad \text{and} \quad \rho^0(\cdot) = \rho(\cdot, 0).$$

Also our calculations frequently use the following exponents

$$a = \frac{\alpha_N}{\alpha_N + 1} = \frac{\deg(g)}{\deg(g) + 1}, \quad (2.1)$$

and

$$\beta = 2 - a, \quad \lambda = \frac{2 - a}{1 - a} = \frac{\beta}{\beta - 1}, \quad \gamma = \frac{a}{2 - a} = \frac{a}{\beta}. \quad (2.2)$$

The arguments $C, C_0, C_1, \dots$ will represent for positive generic constants and their values depend on exponents, coefficients of polynomial g , the spatial dimension d and domain Ω , independent of the initial and boundary data, size of mesh and time step. These constants may be different place by place.

LEMMA 2.1 (cf. [2, 17]). *The function $K(\xi)$ has the following properties*

- (i) $K : [0, \infty) \rightarrow (0, a_0^{-1}]$ and it decreases in ξ ,
- (ii) For any $n \geq 1$, the function $K(\xi)\xi^n$ increasing and $K(\xi)\xi^n \geq 0$
- (iii) Type of degeneracy

$$\frac{c_1}{(1 + \xi)^a} \leq K(\xi) \leq \frac{c_2}{(1 + \xi)^a}, \quad (2.3)$$

(iv) For all $n \geq 1$,

$$c_3(\xi^{n-a} - 1) \leq K(\xi)\xi^n \leq c_2\xi^{n-a}, \quad (2.4)$$

(iv) Relation with its derivative

$$-aK(\xi) \leq K'(\xi)\xi \leq 0, \quad (2.5)$$

where c_1, c_2, c_3 are positive constants depending on Ω and g .

We define

$$H(\xi) = \int_0^{\xi^2} K(\sqrt{s})dx, \text{ for } \xi \geq 0. \quad (2.6)$$

The function $H(\xi)$ can compare with ξ and $K(\xi)$ by

$$K(\xi)\xi^2 \leq H(\xi) \leq 2K(\xi)\xi^2. \quad (2.7)$$

For the monotonicity and continuity of the differential operator in (1.15) we have the following results.

LEMMA 2.2 (cf. [17]). *One has*

(i) For all $y, y' \in \mathbb{R}^d$,

$$(K(|y'|)y' - K(|y|)y) \cdot (y' - y) \geq (\beta - 1)K(\max\{|y|, |y'|\})|y' - y|^2. \quad (2.8)$$

(ii) For the vector functions s_1, s_2 , there is a positive constant C such that

$$(K(|s_1|)s_1 - K(|s_2|)s_2, s_1 - s_2) \geq C\omega \|s_1 - s_2\|_{L^\beta(\Omega)}^2, \quad (2.9)$$

where $\omega = (1 + \max\{\|s_1\|_{L^\beta(\Omega)}, \|s_2\|_{L^\beta(\Omega)}\})^{-a}$.

LEMMA 2.3 (cf. [26]). *For all $y, y' \in \mathbb{R}^d$. There exist a positive constant C depending on polynomial g , the spatial dimension d and domain Ω such that*

$$|K(|y'|)y' - K(|y|)y| \leq C|y' - y|. \quad (2.10)$$

Next we derive trace estimates suitable to our nonlinear problem.

LEMMA 2.4. (i) Let $q \geq 2$. Assume $v(x)$ is a function defined on Ω . If $|v|^{q-1} \in W^{1,1}(\Omega)$ then for all $\varepsilon > 0$,

$$\int_{\Gamma} |v|^{q-1} d\sigma \leq C \|v\|_{L^q(\Omega)}^{q-1} + \varepsilon \int_{\Omega} |v|^{q-2} |\nabla v|^\beta dx + C \varepsilon^{-\frac{1}{\beta-1}} \|v\|_{L^q(\Omega)}^{q-2}. \quad (2.11)$$

(ii) If $u \in L^\infty(\Gamma)$ and $|v| \in W^{1,1}(\Omega)$ then for all $\varepsilon > 0$,

$$|\langle u, v \rangle| \leq \varepsilon \left(\|v\|^2 + \|\nabla v\|_{L^\beta(\Omega)}^\beta \right) + C \left(\varepsilon^{-1} \|u\|_{L^\infty(\Gamma)}^2 + \varepsilon^{-\frac{1}{\beta-1}} \|u\|_{L^\infty(\Gamma)}^\lambda \right). \quad (2.12)$$

In particular case,

$$|\langle u, v \rangle| \leq \frac{1}{4} \left(\|v\|^2 + \|\nabla v\|_{L^\beta(\Omega)}^\beta \right) + C \left(1 + \|u\|_{L^\infty(\Gamma)}^\lambda \right). \quad (2.13)$$

Above, C is a positive constant independent of u, v, ε .

Proof. We recall the trace theorem

$$\int_{\Gamma} |\phi| dx \leq C \int_{\Omega} |\phi| dx + C \int_{\Omega} |\nabla \phi| dx,$$

for all $\phi \in W^{1,1}(\Omega)$, where C are positive constants depending on Ω . Applying the Trace theorem to $\phi = |v|^{q-1}$ shows that

$$\int_{\Gamma} |v|^{q-1} d\sigma \leq C \int_{\Omega} |v|^{q-1} dx + C \int_{\Omega} |v|^{q-2} |\nabla v| dx. \quad (2.14)$$

Note that $\frac{1}{\beta} + \frac{1}{\lambda} = 1$ and $\frac{\lambda}{\beta} = \frac{1}{\beta-1}$. Using Young's inequality with exponent β and λ , we find that for $\varepsilon > 0$

$$\begin{aligned} |v|^{q-2} |\nabla v| &= \varepsilon^{\frac{1}{\beta}} |\rho_h|^{\frac{q-2}{\beta}} |\nabla v| \cdot \varepsilon^{-\frac{1}{\beta}} |v|^{\frac{q-2}{\lambda}} \\ &\leq \varepsilon |v|^{q-2} |\nabla v|^\beta + C \varepsilon^{-\frac{1}{\beta-1}} |v|^{q-2}. \end{aligned} \quad (2.15)$$

Combining (2.14), (2.15) we obtain

$$\int_{\Gamma} |v|^{q-1} d\sigma \leq C \int_{\Omega} |v|^{q-1} dx + \varepsilon \int_{\Omega} |v|^{q-2} |\nabla v|^\beta dx + C \varepsilon^{-\frac{1}{\beta-1}} \int_{\Omega} |v|^{q-2} dx. \quad (2.16)$$

Inequality (2.11) follows by using Hölder's inequality to the first term of the right hand side in (2.16).

(ii) We have

$$|\langle u, v \rangle| \leq \|u\|_{L^\infty(\Gamma)} \int_{\Gamma} |v| d\sigma. \quad (2.17)$$

For all $\delta > 0$, using (2.16) with $q = 2$ gives,

$$\int_{\Gamma} |v| d\sigma \leq C \int_{\Omega} |v| dx + \delta \|\nabla v\|_{L^\beta}^\beta + C \delta^{-\frac{1}{\beta-1}}.$$

Applying Young's inequality leads to

$$\int_{\Gamma} |v| d\sigma \leq \delta \|v\|^2 + C \delta^{-1} + \delta \|\nabla v\|_{L^\beta}^\beta + C \delta^{-\frac{1}{\beta-1}},$$

which gives

$$|\langle u, v \rangle| \leq \|u\|_{L^\infty(\Gamma)} \left(\delta \|v\|^2 + C \delta^{-1} + \delta \|\nabla v\|_{L^\beta}^\beta + C \delta^{-\frac{1}{\beta-1}} \right).$$

If $\|u\|_{L^\infty(\Gamma)} = 0$ then (2.12) clearly holds true.

Otherwise, selecting $\delta = \varepsilon \|u\|_{L^\infty(\Gamma)}^{-1}$ and the fact that $\frac{1}{\beta-1} + 1 = \lambda$ we obtain estimate (2.12).

Estimate (2.13) follows by choosing $\varepsilon = 1/4$ in (2.12) and using Young's inequality. $\square$

3. The Galerkin finite element method. Our aim is to study equation (1.15) for density of slightly compressible fluids in bounded domain in porous media. The fluid flows are subject to some conditions on the boundary.

We consider the initial boundary value problem associated with (1.15) ,

$$\begin{cases} \rho_t - \nabla \cdot (K(|\nabla \rho|)\nabla \rho) = f & \text{in } \Omega \times I, \\ \rho(x, 0) = \rho^0(x) & \text{in } \Omega, \\ K(|\nabla \rho|)\nabla \rho \cdot \nu + \psi = 0 & \text{on } \Gamma \times I, \end{cases} \quad (3.1)$$

where $\rho^0(x)$ and $\psi(x, t)$ are given initial and boundary data, respectively.

The variational formulation of (3.1) is defined as the following:

Find $\rho : [0, T] \rightarrow W \equiv H^1(\Omega)$ such that

$$(\rho_t, w) + (K(|\nabla \rho|)\nabla \rho, \nabla w) = -\langle \psi, w \rangle + (f, w), \quad w \in W \quad (3.2)$$

with $\rho(x, 0) = \rho^0(x)$.

Let $\{\mathcal{T}_h\}_h$ be a family of quasi-uniform triangulations of Ω with h being the maximum diameter of the element. Let W_h the space of discontinuous piecewise polynomials of degree $r \geq 0$ over $\mathcal{T}_h$. It is frequently valuable to decompose the analysis of the convergence of finite element methods by passing through a projection of the solution of the differential problem into the finite element space. Here we use the standard L^2 -projection operator (see [6]) $\pi : W \rightarrow W_h$, satisfying

$$(\pi w, v_h) = (w, v_h), \quad \forall w \in W, v_h \in W_h.$$

These projections have well-known approximation properties (see [5, 25]).

(i) $\|\pi w\| \leq \|w\|$ holds for all $w \in L^2(\Omega)$.

(ii) There exist positive constants C_1, C_2 such that

$$\|\pi w - w\|_{L^q(\Omega)} \leq C_1 h^m \|w\|_{W^{m,q}(\Omega)} \quad (3.3)$$

for all $w \in W^{m,q}(\Omega)$, $0 \leq m \leq r+1$, $1 \leq q \leq \infty$. Notation $\|\cdot\|_{m,q}$ denotes a standard norm in Sobolev space $W^{m,q}(\Omega)$. In short hand, when $q = 2$ we write (3.3) as

$$\|\pi w - w\| \leq C_1 h^m \|w\|_m \quad \text{and} \quad \|\pi w - w\|_{L^\infty(\Omega)} \leq C_2 h^m \|w\|_{m+1}.$$

Replacing the original density by its approximation, the semidiscrete formulation of (3.2) can read as following: Find $\rho_h : [0, T] \rightarrow W_h$ such that

$$(\rho_{h,t}, w_h) + (K(|\nabla \rho_h|)\nabla \rho_h, \nabla w_h) = -\langle \psi, w_h \rangle + (f, w_h), \quad w_h \in W_h \quad (3.4)$$

with initial data $\rho_h^0 = \pi \rho^0(x)$.

Let N be the positive integer, $t_0 = 0 < t_1 < \dots < t_N = T$ be partition interval $[0, T]$ of N sub-intervals, and let $\Delta t = t_n - t_{n-1} = T/N$ be the n -th time step size, $t_n = n\Delta t$ and $\varphi^n = \varphi(\cdot, t_n)$. The discrete time Galerkin finite element approximation to (3.2) is defined as follows:

Find $\rho_h^n \in W_h$, $n = 1, 2, \dots, N$, such that

$$\left(\frac{\rho_h^n - \rho_h^{n-1}}{\Delta t}, w_h \right) + (K(|\nabla \rho_h^n|)\nabla \rho_h^n, \nabla w_h) = -\langle \psi^n, w_h \rangle + (f^n, w_h), \quad \forall w_h \in W_h, \quad (3.5)$$

with initial data are chosen as follows: $\rho_h^0(x) = \pi \rho^0(x)$.

4. A priori estimate for solutions. We study the (3.2), (3.4) and (3.5) equations for the density with fixed the functions $g(s)$ in (1.7) and (1.8). Therefore, the exponents α_i and coefficients a_i are all fixed, and so are the function $K(\xi)$, $H(\xi)$ in (1.12), (2.6).

With properties (2.3), (2.4), (2.5), the monotonicity (2.8), and by classical theory of monotone operators [28, 33, 38], the authors in [19] proved the global existence and uniqueness of weak solution of equation (3.2). Moreover $\rho \in C(0, T, L^q(\Omega)) \cap L_{loc}^\beta(0, T, W^{1,\beta}(\Omega))$, $q \geq 1$ and $\rho_t \in L_{loc}^{\beta'}(0, T, (W^{1,\beta}(\Omega))') \cap L_{loc}^2(0, T, L^2(\Omega))$ provided the initial, boundary data and f sufficiently smooth. For a *a priori* estimate, we assume throughout this paper that ψ, ψ_t belong to $C([0, T], L^\infty(\Gamma))$, $\rho^0 \in L^\infty(\Omega)$ and $f \in C([0, T], L^\infty(\Omega))$ and the weak solution are sufficiently regularities both in x and t variables such that our calculations can be performed legitimately.

The local existence of approximate solution of (3.4) follows from Peano's theorem. The global existence of ρ_h relies on the known theory of ordinary differential equation and stability estimate which shows as follows.

THEOREM 4.1. *Let $q \geq 2$ and ρ_h be a solution of (3.4). Then there exist positive constant C such that*

$$\|\rho_h\|_{L^\infty(I, L^q(\Omega))} \leq C \|\rho^0\|_{L^q(\Omega)} + C \left[\int_0^T \left(1 + \|\psi\|_{L^\infty(\Gamma)}^{\frac{q\lambda}{2}} + \|f\|_{L^q(\Omega)}^q \right) dt \right]^{\frac{1}{q}}. \quad (4.1)$$

Proof. Selecting $w_h = |\rho_h|^{q-1} \text{sign}(\rho_h)$ in (3.4), we obtain

$$\begin{aligned} \frac{1}{q} \frac{d}{dt} \|\rho_h\|_{L^q}^q &= -(q-1) (K(|\nabla \rho_h|) |\nabla \rho_h|^2, |\rho_h|^{q-2}) \\ &\quad - \langle \psi, |\rho_h|^{q-1} \text{sign}(\rho_h) \rangle + (f, |\rho_h|^{q-1} \text{sign}(\rho_h)). \end{aligned} \quad (4.2)$$

It follows from (2.4) and Hölder's inequality that

$$\begin{aligned} -(q-1) (K(|\nabla \rho_h|) |\nabla \rho_h|^2, |\rho_h|^{q-2}) &\leq -c_0 \int_{\Omega} (|\nabla \rho_h|^\beta - 1) |\rho_h|^{q-2} dx \\ &= -c_0 \int_{\Omega} |\nabla \rho_h|^\beta |\rho_h|^{q-2} dx + c_0 \int_{\Omega} |\rho_h|^{q-2} dx \\ &\leq -c_0 \int_{\Omega} |\nabla \rho_h|^\beta |\rho_h|^{q-2} dx + C \|\rho_h\|_{L^q}^{q-2}, \end{aligned} \quad (4.3)$$

where $c_0 = c_3(q-1)$.

According to (2.11) in Lemma 2.4,

$$\begin{aligned} |-\langle \psi, |\rho_h|^{q-1} \text{sign}(\rho_h) \rangle| &\leq \|\psi\|_{L^\infty(\Gamma)} \int_{\Gamma} |\rho_h|^{q-1} d\sigma \\ &\leq C \|\psi\|_{L^\infty(\Gamma)} \left\{ \|\rho_h\|_{L^q}^{q-1} + \varepsilon \int_{\Omega} |\rho_h|^{q-2} |\nabla \rho_h|^\beta dx + \varepsilon^{-\frac{1}{\beta-1}} \|\rho_h\|_{L^q}^{q-2} \right\}. \end{aligned} \quad (4.4)$$

Using Young's inequalities with exponent $q/(q-1)$ and q , we obtain

$$(f, |\rho_h|^{q-1} \text{sign}(\rho_h)) \leq C \|\rho_h\|_{L^q}^{q-1} + C \|f\|_{L^q}^q. \quad (4.5)$$

Combining (4.3), (4.4), (4.5) and (4.2) gives

$$\begin{aligned} \frac{1}{q} \frac{d}{dt} \|\rho_h\|_{L^q}^q &\leq -c_0 \int_{\Omega} |\nabla \rho_h|^\beta |\rho_h|^{q-2} dx + C \|\rho_h\|_{L^q}^{q-2} \\ &\quad + C \|\psi\|_{L^\infty(\Gamma)} \left\{ \|\rho_h\|_{L^q}^{q-1} + \varepsilon \int_{\Omega} |\rho_h|^{q-2} |\nabla \rho_h|^\beta dx + \varepsilon^{-\frac{1}{\beta-1}} \|\rho_h\|_{L^q}^{q-2} \right\} \\ &\quad + C \|\rho_h\|_{L^q}^{q-1} + C \|f\|_{L^q}^q. \end{aligned} \quad (4.6)$$

The case $\|\psi(t)\|_{L^\infty(\Gamma)} = 0$, by the means of Young's inequality in (4.6) we obtain

$$\frac{d}{dt} \|\rho_h\|_{L^q}^q + qc_0 \int_{\Omega} |\nabla \rho_h|^\beta |\rho_h|^{q-2} dx \leq C \|\rho_h\|_{L^q}^q + C(1 + \|f\|_{L^q}^q). \quad (4.7)$$

Applying Gronwall's inequality to above differential inequality implies that.

$$\begin{aligned} \|\rho_h\|_{L^\infty(I, L^q(\Omega))}^q + qc_0 \int_0^t \int_{\Omega} e^{C(t-\tau)} |\nabla \rho_h|^\beta |\rho_h|^{q-2} dx d\tau \\ \leq C \|\rho_h^0\|_{L^q(\Omega)}^q + C \int_0^T \left(1 + \|f\|_{L^q(\Omega)}^q\right) dt. \end{aligned} \quad (4.8)$$

Consider $\|\psi(t)\|_{L^\infty(\Gamma)} \neq 0$. Selecting $\varepsilon = \frac{c_0}{2C} \|\psi\|_{L^\infty(\Gamma)}^{-1}$ in (4.6), we obtain

$$\begin{aligned} \frac{d}{dt} \|\rho_h\|_{L^q}^q \leq -\frac{qc_0}{2} \int_{\Omega} |\nabla \rho_h|^\beta |\rho_h|^{q-2} dx + C \|\rho_h\|_{L^q}^{q-2} \\ + C \|\psi\|_{L^\infty(\Gamma)} \left\{ \|\rho_h\|_{L^q}^{q-1} + \|\psi\|_{L^\infty(\Gamma)}^{\frac{1}{\beta-1}} \|\rho_h\|_{L^q}^{q-2} \right\} + C \|\rho_h\|_{L^q}^{q-1} + C \|f\|_{L^q}^q. \end{aligned} \quad (4.9)$$

By Young's inequality,

$$\begin{aligned} \frac{d}{dt} \|\rho_h\|_{L^q}^q + \frac{qc_0}{2} \int_{\Omega} |\nabla \rho_h|^\beta |\rho_h|^{q-2} dx \\ \leq C \|\rho_h\|_{L^q}^q + C \left(1 + \|\psi\|_{L^\infty(\Gamma)}^q + \|\psi\|_{L^\infty(\Gamma)}^{\frac{q\beta}{2(\beta-1)}} + \|f\|_{L^q}^q\right) \\ \leq C \|\rho_h\|_{L^q}^q + C \left(1 + \|\psi\|_{L^\infty(\Gamma)}^{\frac{q\beta}{2(\beta-1)}} + \|f\|_{L^q}^q\right). \end{aligned} \quad (4.10)$$

Solving this differential inequality shows that,

$$\begin{aligned} \|\rho_h\|_{L^q}^q + \frac{qc_0}{2} \int_0^t \int_{\Omega} e^{C(t-\tau)} |\nabla \rho_h|^\beta |\rho_h|^{q-2} dx d\tau \\ \leq \|\rho_h^0\|_{L^q}^q + C \int_0^t \left(1 + \|\psi\|_{L^\infty(\Gamma)}^{\frac{q\beta}{2(\beta-1)}} + \|f\|_{L^q}^q\right) d\tau. \end{aligned} \quad (4.11)$$

It is easy to see that in both above cases

$$\|\rho_h\|_{L^q}^q \leq \|\rho_h^0\|_{L^q}^q + C \int_0^t \left(1 + \|\psi\|_{L^\infty(\Gamma)}^{\frac{q\beta}{2(\beta-1)}} + \|f\|_{L^q}^q\right) d\tau.$$

Using inequality $(a+b)^{1/q} \leq 2^{1/q}(a^{1/q} + b^{1/q})$ with

$$a = \|\rho_h^0\|_{L^q}^q \quad \text{and} \quad b = \int_0^t \left(1 + \|\psi\|_{L^\infty(\Gamma)}^{\frac{q\beta}{2(\beta-1)}} + \|f\|_{L^q}^q\right) d\tau,$$

we obtain

$$\|\rho_h\|_{L^\infty(I, L^q(\Omega))} \leq C \|\rho_h^0\|_{L^q(\Omega)} + C \left[\int_0^T \left(1 + \|\psi\|_{L^\infty(\Gamma)}^{\frac{q\beta}{2(\beta-1)}} + \|f\|_{L^q(\Omega)}^q\right) dt \right]^{\frac{1}{q}}. \quad (4.12)$$

Note that

$$\|\rho_h^0\| = \|\pi \rho^0\| \leq \|\rho^0\|. \quad (4.13)$$

Thus inequality (4.1) holds. We finish the proof. $\square$

Next we give estimates in L^2 -norm following directly from (4.11) and (4.13) with $q = 2$.

LEMMA 4.2. *Under assumption of Theorem 4.1. There exist a positive constant C such that*

$$\begin{aligned} & \|\rho_h\|_{L^\infty(I, L^2(\Omega))} + \|\nabla \rho_h\|_{L^\beta(I, L^\beta(\Omega))}^{\frac{\beta}{2}} \\ & \leq C \|\rho^0\| + C \left[\int_0^T \left(1 + \|\psi(t)\|_{L^\infty(\Gamma)}^\lambda + \|f(t)\|^2 \right) dt \right]^{\frac{1}{2}}. \end{aligned} \quad (4.14)$$

Since the equation (3.4) can interpret as the finite system of ordinary differential equations in the coefficients of ρ_h with respect to basis of W_h . The stability estimate (4.12) suffices to establish existence of $\rho_h(t)$ for all $t \in I$.

For the uniqueness of approximation solution, assume that for $i = 1, 2$, $\rho_h^{(i)} \in W_h$ is the solution of (3.4). Let $\rho_h = \rho_h^{(1)} - \rho_h^{(2)}$ then

$$(\rho_{h,t}, w_h) + (K(|\nabla \rho_h^{(1)}|) \nabla \rho_h^{(1)} - K(|\nabla \rho_h^{(2)}|) \nabla \rho_h^{(2)}, \nabla w_h) = 0.$$

Choose $w_h = \rho_h$, implies

$$\frac{1}{2} \frac{d}{dt} \|\rho_h\|^2 + \left(K(|\nabla \rho_h^{(1)}|) \nabla \rho_h^{(1)} - K(|\nabla \rho_h^{(2)}|) \nabla \rho_h^{(2)}, \nabla \rho_h^{(1)} - \nabla \rho_h^{(2)} \right) = 0.$$

According to (2.9) we find that

$$\begin{aligned} & \left(K(|\nabla \rho_h^{(1)}|) \nabla \rho_h^{(1)} - K(|\nabla \rho_h^{(2)}|) \nabla \rho_h^{(2)}, \nabla \rho_h^{(1)} - \nabla \rho_h^{(2)} \right) \\ & \geq C \left[1 + \max\{\|\nabla \rho_h^{(1)}\|_{L^\beta}, \|\nabla \rho_h^{(2)}\|_{L^\beta}\} \right]^{-a} \|\nabla \rho_h\|_{L^\beta}^2. \end{aligned}$$

Hence

$$\frac{1}{2} \frac{d}{dt} \|\rho_h\|^2 + C \left[1 + \max\{\|\nabla \rho_h^{(1)}\|_{L^\beta}, \|\nabla \rho_h^{(2)}\|_{L^\beta}\} \right]^{-a} \|\nabla \rho_h\|_{L^\beta}^2 \leq 0.$$

Integrating in time from 0 to t we have

$$\|\rho_h\|^2 + C \int_0^t \left[1 + \max\{\|\nabla \rho_h^{(1)}\|_{L^\beta}, \|\nabla \rho_h^{(2)}\|_{L^\beta}\} \right]^{-a} \|\nabla \rho_h\|_{L^\beta}^2 dt \leq \|\rho_h(0)\|^2 = 0.$$

Thus

$$\rho_h = 0, \quad \nabla \rho_h = 0 \quad \forall (x, t) \in \Omega \times I.$$

Next, we find estimates for $\nabla \rho_h$. We define

$$g(t) = 1 + \|\psi(t)\|_{L^\infty(\Gamma)}^\lambda, \quad h(t) = 1 + \|\psi_t(t)\|_{L^\infty(\Gamma)}^\lambda, \quad k(t) = g(t) + \|f(t)\|^2. \quad (4.15)$$

THEOREM 4.3. *Let ρ_h be a solution to the problem (3.4). Then there exist constant positive constant C satisfying*

$$\|\rho_{h,t}\|_{L^2(I, L^2(\Omega))}^2 + \|\nabla \rho_h\|_{L^\beta(I, L^\beta(\Omega))}^\beta + \|\nabla \rho_h\|_{L^\infty(I, L^\beta(\Omega))}^\beta + \|\rho_h\|_{L^\infty(I, L^2(\Omega))}^2 \leq C\mathcal{M}. \quad (4.16)$$

where

$$\begin{aligned} \mathcal{M} = & \max_{t \in [0, T]} g(t) + \int_0^T [h(t) + k(t)] dt + \int_0^T \int_0^t k(\tau) d\tau dt \\ & + \|\rho^0\|^2 + \|\nabla \rho^0\|_{L^\beta(\Omega)}^\beta + \|\psi^0\|_{L^2(\Gamma)} \|\rho^0\|_{L^2(\Gamma)}. \end{aligned} \quad (4.17)$$

Proof. Choosing $w_h = \rho_{h,t}$ in (3.4) leads to

$$\begin{aligned} \|\rho_{h,t}\|^2 + \frac{1}{2} \frac{d}{dt} \int_{\Omega} H(x, t) dx &= -\langle \psi, \rho_{h,t} \rangle + (f, \rho_{h,t}) \\ &= -\frac{d}{dt} \langle \psi, \rho_h \rangle + \langle \psi_t, \rho_h \rangle + (f, \rho_{h,t}). \end{aligned} \quad (4.18)$$

where $H(x, t) = H(|\nabla \rho_h(x, t)|)$ is defined in (2.6).

With $q = 2$ then from (4.10),

$$\frac{d}{dt} \|\rho_h\|^2 + c_0 \|\nabla \rho_h\|_{L^\beta}^\beta \leq C \|\rho_h\|^2 + C \left(1 + \|\psi\|_{L^\infty(\Gamma)}^\lambda + \|f\|^2 \right). \quad (4.19)$$

Let

$$\mathcal{E}(t) = \int_{\Omega} H(x, t) dx + \|\rho_h\|^2 + 2 \langle \psi, \rho_h \rangle.$$

Adding two inequalities (4.18) and (4.19) gives

$$\|\rho_{h,t}\|^2 + c_0 \|\nabla \rho_h\|_{L^\beta}^\beta + \frac{1}{2} \frac{d}{dt} \mathcal{E}(t) \leq \langle \psi_t, \rho_h \rangle + (f, \rho_{h,t}) + C \|\rho_h\|^2 + Ck(t).$$

Using (2.13) and Cauchy's inequality then

$$\begin{aligned} \|\rho_{h,t}\|^2 + c_0 \|\nabla \rho_h\|_{L^\beta}^\beta + \frac{1}{2} \frac{d}{dt} \mathcal{E}(t) &\leq \frac{1}{2} \left(\|\rho_h\|^2 + \|\nabla \rho_h\|_{L^\beta}^\beta \right) + C \left(1 + \|\psi_t\|_{L^\infty(\Gamma)}^\lambda \right) \\ &\quad + \frac{1}{2} \|f\|^2 + \frac{1}{2} \|\rho_{h,t}\|^2 + C \|\rho_h\|^2 + Ck(t). \end{aligned}$$

Absorbing $\|f\|^2$ to $k(t)$, integrating in time, we obtain

$$\begin{aligned} \int_0^T \|\rho_{h,t}\|^2 dt + c_0 \int_0^T \|\nabla \rho_h\|_{L^\beta}^\beta dt + \int_{\Omega} H(x, t) dx + \|\rho_h\|^2 &\leq -2 \langle \psi, \rho_h \rangle \\ &\quad + C \int_0^T \|\rho_h\|^2 dt + C \int_0^T [h(t) + k(t)] dt + \mathcal{E}(0). \end{aligned} \quad (4.20)$$

Applying (2.12) to the first term of the left hand side in (4.20) and using the fact in (2.7) that $c_3(|\nabla \rho_h|^\beta - 1) \leq H(x, t) \leq 2c_2|\nabla \rho_h|^\beta$, we have

$$\begin{aligned} \int_0^T \|\rho_{h,t}\|^2 dt + c_0 \int_0^T \|\nabla \rho_h\|_{L^\beta}^\beta dt + c_3 \|\nabla \rho_h\|_{L^\beta}^\beta + \|\rho_h\|^2 \\ \leq 2\varepsilon \left(\|\rho_h\|^2 + \|\nabla \rho_h\|_{L^\beta}^\beta \right) + C \left(\varepsilon^{-1} \|\psi\|_{L^\infty(\Gamma)} + \varepsilon^{-\frac{1}{\beta-1}} \|\psi\|_{L^\infty(\Gamma)}^\lambda \right) \\ + C \int_0^T \|\rho_h\|^2 dt + C \int_0^T [h(t) + k(t)] dt + \mathcal{E}(0). \end{aligned} \quad (4.21)$$

Thus by taking $\varepsilon = \min\{c_3, 1\}/4$, and using Young's inequality,

$$\begin{aligned} \int_0^T \|\rho_{h,t}\|^2 dt + c_0 \int_0^T \|\nabla \rho_h\|_{L^\beta}^\beta dt + \frac{c_3}{2} \|\nabla \rho_h\|_{L^\beta}^\beta + \frac{1}{2} \|\rho_h\|^2 &\leq Cg(t) \\ + C \int_0^T \|\rho_h\|^2 dt + C \int_0^T [h(t) + k(t)] dt + \mathcal{E}(0). \end{aligned} \quad (4.22)$$

Note from (4.11) with $q = 2$ that

$$\begin{aligned} \int_0^T \|\rho_h\|^2 dt &\leq C \left(T \|\rho_h(0)\|^2 + \int_0^T \int_0^t k(\tau) d\tau dt \right) \\ &\leq C \left(T \|\rho^0\|^2 + \int_0^T \int_0^t k(\tau) d\tau dt \right) \end{aligned}$$

and from (2.7), (2.4) that

$$\begin{aligned}\mathcal{E}(0) &\leq C \left(\|\rho_h^0\|^2 + \|\nabla \rho_h^0\|_{L^\beta(\Omega)}^\beta + \|\psi(0)\|_{L^2(\Gamma)} \|\rho_h^0\|_{L^2(\Gamma)} \right) \\ &\leq C \left(\|\rho^0\|^2 + \|\nabla \rho^0\|_{L^\beta(\Omega)}^\beta + \|\psi(0)\|_{L^2(\Gamma)} \|\rho^0\|_{L^2(\Gamma)} \right).\end{aligned}$$

The left hand side of (4.22) is bounded by $C\mathcal{M}$ which implies (4.16). $\square$

Now we prove that the time derivative of density is also bounded.

THEOREM 4.4. *Let $0 < t_0 < T$, ρ_h be a solution to the semidiscrete problem (3.4). Then there exist constant positive constant C such that*

$$\|\rho_{h,t}\|_{L^\infty(I, L^2(\Omega))}^2 \leq C t_0^{-1} \mathcal{M} + C \mathcal{M} + C \int_0^T h(t) \left(1 + \|f_t(t)\|^2 \right) dt. \quad (4.23)$$

where $\mathcal{M}$ and $h(t)$ are defined in (4.17) and (4.15) respectively.

Proof. Differentiating (3.4) with respect t yields that

$$\begin{aligned}(\rho_{h,tt}, w_h) + (K(|\nabla \rho_h|) \nabla \rho_{h,t}, \nabla w_h) \\ = - \left(K'(|\nabla \rho_h|) \frac{\nabla \rho_h \cdot \nabla \rho_{h,t}}{|\nabla \rho_h|} \nabla \rho_h, \nabla w_h \right) - \langle \psi_t, w_h \rangle + (f_t, w_h).\end{aligned}$$

Choosing $w_h = \rho_{h,t}$, we obtain

$$\begin{aligned}\frac{1}{2} \frac{d}{dt} \|\rho_{h,t}\|^2 + \left\| K^{1/2}(|\nabla \rho_h|) \nabla \rho_{h,t} \right\|^2 \\ = - \left(K'(|\nabla \rho_h|) \frac{\nabla \rho_h \cdot \nabla \rho_{h,t}}{|\nabla \rho_h|} \nabla \rho_h, \nabla \rho_{h,t} \right) + (f_t, \rho_{h,t}) - \langle \psi_t, \rho_{h,t} \rangle.\end{aligned}$$

Using (2.5),

$$\left| - \left(K'(|\nabla \rho_h|) \frac{\nabla \rho_h \cdot \nabla \rho_{h,t}}{|\nabla \rho_h|} \nabla \rho_h, \nabla \rho_{h,t} \right) \right| \leq a \left\| K^{1/2}(|\nabla \rho_h|) \nabla \rho_{h,t} \right\|^2. \quad (4.24)$$

Thus

$$\frac{1}{2} \frac{d}{dt} \|\rho_{h,t}\|^2 + (1-a) \left\| K^{1/2}(|\nabla \rho_h|) \nabla \rho_{h,t} \right\|^2 \leq (f_t, \rho_{h,t}) - \langle \psi_t, \rho_{h,t} \rangle. \quad (4.25)$$

In virtue of Cauchy's inequality, for all $\varepsilon > 0$

$$(f_t, \rho_{h,t}) \leq \varepsilon \|\rho_{h,t}\|^2 + C \varepsilon^{-1} \|f_t\|^2. \quad (4.26)$$

Using Trace Theorem we obtain,

$$|\langle \psi_t, \rho_{h,t} \rangle| \leq \|\psi_t\|_{L^\infty(\Gamma)} [(\rho_{h,t}, 1) + (|\nabla \rho_{h,t}|, 1)]. \quad (4.27)$$

Again Cauchy's inequality gives that for all $\varepsilon, \varepsilon_1 > 0$

$$(\rho_{h,t}, 1) \leq \varepsilon \|\rho_{h,t}\|^2 + C \varepsilon^{-1}, \quad (4.28)$$

and

$$(|\nabla \rho_{h,t}|, 1) \leq \varepsilon_1 (K(|\nabla \rho_h|) |\nabla \rho_{h,t}|^2, 1) + C \varepsilon_1^{-1} (K^{-1}(|\nabla \rho_h|), 1).$$

By using (2.3) and $(1+x)^a \leq 1+x^a, x \geq 0$ imply

$$\begin{aligned}(|\nabla \rho_{h,t}|, 1) &\leq \varepsilon_1 (K(|\nabla \rho_h|) |\nabla \rho_{h,t}|^2, 1) + C \varepsilon_1^{-1} (1 + |\nabla \rho_h|^a, 1) \\ &\leq \varepsilon_1 \left\| K^{1/2}(|\nabla \rho_h|) \nabla \rho_{h,t} \right\|^2 + C \varepsilon_1^{-1} (1 + \|\nabla \rho_h\|_{L^\beta}^a).\end{aligned} \quad (4.29)$$

Combining (4.27), (4.28) and (4.29) shows that

$$|\langle \psi_t, \rho_{h,t} \rangle| \leq \|\psi_t\|_{L^\infty(\Gamma)} \left\{ \varepsilon \|\rho_{h,t}\|^2 + C\varepsilon^{-1} + \varepsilon_1 \left\| K^{1/2}(|\nabla \rho_h|) \nabla \rho_{h,t} \right\|^2 + C\varepsilon_1^{-1} (1 + \|\nabla \rho_h\|_{L^\beta}^a) \right\}. \quad (4.30)$$

It follows from (4.25), (4.26) and (4.30) that

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\rho_{h,t}\|^2 + (1-a) \left\| K^{1/2}(|\nabla \rho_h|) \nabla \rho_{h,t} \right\|^2 &\leq \|\psi_t\|_{L^\infty(\Gamma)} \left\{ \varepsilon \|\rho_{h,t}\|^2 + C\varepsilon^{-1} \right. \\ &\quad \left. + \varepsilon_1 \left\| K^{1/2}(|\nabla \rho_h|) \nabla \rho_{h,t} \right\|^2 + C\varepsilon_1^{-1} (1 + \|\nabla \rho_h\|_{L^\beta}^a) \right\} + \varepsilon \|\rho_{h,t}\|^2 + C\varepsilon^{-1} \|f_t\|^2. \end{aligned} \quad (4.31)$$

If $\|\psi_t(t)\|_{L^\infty(\Gamma)} = 0$ then

$$\frac{1}{2} \frac{d}{dt} \|\rho_{h,t}\|^2 + (1-a) \left\| K^{1/2}(|\nabla \rho_h|) \nabla \rho_{h,t} \right\|^2 \leq \varepsilon \|\rho_{h,t}\|^2 + C\varepsilon^{-1} \|f_t\|^2.$$

Dropping the second term of the left hand side in above estimate, selecting $\varepsilon = 1/2$ and, for $0 < t' \leq t_0 < t \leq T$ integrating from t' to t give

$$\begin{aligned} \|\rho_{h,t}(t)\|^2 d\tau &\leq \|\rho_{h,t}(t')\|^2 + \int_{t'}^t \|\rho_{h,t}\|^2 d\tau + C \int_{t'}^t \|f_t\|^2 d\tau \\ &\leq \|\rho_{h,t}(t')\|^2 + \int_0^t \|\rho_{h,t}\|^2 d\tau + C \int_0^t \|f_t\|^2 d\tau. \end{aligned} \quad (4.32)$$

Now integrating in t' from 0 to t_0 ,

$$t_0 \|\rho_{h,t}(t)\|^2 d\tau \leq \int_0^{t_0} \|\rho_{h,t}(t')\|^2 dt' + t_0 \left(\int_0^t \|\rho_{h,t}\|^2 d\tau + C \int_0^t \|f_t\|^2 d\tau \right). \quad (4.33)$$

Using (4.16), we find that

$$t_0 \|\rho_{h,t}(t)\|^2 d\tau \leq \mathcal{M} + Ct_0 \left(\mathcal{M} + \int_0^t \|f_t\|^2 d\tau \right), \quad (4.34)$$

which holds (4.23).

Consider $\|\psi_t\|_{L^\infty(\Gamma)} \neq 0$. Selecting $\varepsilon_1 = \frac{1-a}{2} \|\psi_t\|_{L^\infty(\Gamma)}^{-1}$ and $\varepsilon = \frac{1}{2} (\|\psi_t\|_{L^\infty(\Gamma)} + 1)^{-1}$ in (4.31) yields

$$\begin{aligned} \frac{d}{dt} \|\rho_{h,t}\|^2 + (1-a) \|K(|\nabla \rho_h|) \nabla \rho_{h,t}\|^2 &\leq \|\rho_{h,t}\|^2 + C \|\psi_t\|_{L^\infty(\Gamma)}^2 \|\nabla \rho_h\|_{L^\beta}^a \\ &\quad + C \left(\|\psi_t\|_{L^\infty(\Gamma)} + 1 \right) \left(\|\psi_t\|_{L^\infty(\Gamma)} + \|f_t\|^2 \right) + C \|\psi_t\|_{L^\infty(\Gamma)}^2 \\ &\leq \|\rho_{h,t}\|^2 + \|\nabla \rho_h\|_{L^\beta}^\beta + CZ(t), \end{aligned}$$

where

$$Z(t) = \left(\|\psi_t\|_{L^\infty(\Gamma)} + 1 \right) \left(\|\psi_t\|_{L^\infty(\Gamma)} + \|f_t\|^2 \right) + \|\psi_t\|_{L^\infty(\Gamma)}^2 + \|\psi_t\|_{L^\infty(\Gamma)}^\lambda.$$

For $t \geq t_0 \geq t' > 0$. Ignoring the the nonnegative term in the left hand side of above inequality, integrating from t' to t and then integrating in t' from 0 to t_0 , we find that

$$t_0 \|\rho_{h,t}\|^2 \leq \int_0^{t_0} \|\rho_{h,t}(t')\|^2 dt' + t_0 \int_0^t \left[\|\rho_{h,t}\|^2 + \|\nabla \rho_h\|_{L^\beta}^\beta \right] d\tau + Ct_0 \int_0^t Z(\tau) d\tau.$$

By virtue of (4.16),

$$\int_0^t \left[\|\rho_{h,t}\|^2 + \|\nabla \rho_h\|_{L^\beta}^\beta \right] dt \leq C\mathcal{M}, \quad \int_0^{t_0} \|\rho_{h,t}(t')\|^2 dt' \leq C\mathcal{M}.$$

Therefore

$$t_0 \|\rho_{h,t}\|^2 \leq C\mathcal{M} + C\mathcal{M}t_0 + Ct_0 \int_0^t Z(\tau) d\tau. \quad (4.35)$$

We estimate Z -term by

$$\begin{aligned} Z(t) &\leq \left(\|\psi_t\|_{L^\infty(\Gamma)} + 1 \right) \|f_t\|^2 + C \left(1 + \|\psi_t\|_{L^\infty(\Gamma)}^\lambda \right) \\ &\leq C \left(1 + \|\psi_t\|_{L^\infty(\Gamma)}^\lambda \right) \left(1 + \|f_t\|^2 \right). \end{aligned} \quad (4.36)$$

The inequality (4.23) follows from (4.35) and (4.36). The proof is complete. $\square$

The following results can be proved by following the ideas of the proof of Theorem 4.1, 4.3 and 4.4.

THEOREM 4.5. *Let $0 < t_0 < T$, and $q \geq 2$, ρ be a solution to problem (3.2). Then there exist constant positive constant C such that*

$$\|\rho\|_{L^\infty(I, L^q(\Omega))} \leq C \|\rho^0\|_{L^q(\Omega)} + C \left(\int_0^T [1 + \|\psi\|_{L^\infty(\Gamma)}^{\frac{q\lambda}{2}} + \|f(t)\|_{L^q(\Omega)}^q] dt \right)^{\frac{1}{q}}, \quad (4.37)$$

$$\|\rho_t\|_{L^2(I, L^2(\Omega))}^2 + \|\nabla \rho\|_{L^\beta(I, L^\beta(\Omega))}^\beta + \|\nabla \rho\|_{L^\infty(I, L^\beta(\Omega))}^\beta + \|\rho_h\|_{L^\infty(I, L^2(\Omega))}^2 \leq C\mathcal{M}, \quad (4.38)$$

$$\|\rho_t\|_{L^\infty(I, L^2(\Omega))}^2 \leq Ct_0^{-1}\mathcal{M} + C\mathcal{M} + C \int_0^T h(t) \left(1 + \|f(t)\|^2 \right) dt, \quad (4.39)$$

where h and $\mathcal{M}$ are defined as in (4.15) and (4.17).

5. Error estimates. In this section we will establish the error estimates between analytical solution and approximation solution in several norms.

5.1. Error estimate for continuous Galerkin method. We will find the error bounds in the semidiscrete method by comparing the computed solution to the projections of the true solutions. To do this, we restrict the test functions in (3.2) to the finite dimensional spaces. Let

$$\chi = \rho - \rho_h = (\rho - \pi\rho) - (\rho_h - \pi\rho) \stackrel{\text{def}}{=} \vartheta - \theta_h. \quad (5.1)$$

THEOREM 5.1. *Let ρ, ρ_h be two solution to problems (3.2) and (3.4) respectively. Assume that $\rho \in L^\infty(I, H^r(\Omega) \cap W^{r,\beta}(\Omega))$, $\rho_t \in L^2(I, H^r(\Omega))$. Then there exist positive constants C independence of h such that*

$$\begin{aligned} \|\rho - \rho_h\|_{L^\infty(I, L^2(\Omega))} + \mathcal{M}^{-\frac{\gamma}{2}} \|\nabla(\rho - \rho_h)\|_{L^2(I, L^\beta(\Omega))} &\leq Ch^r \|\rho\|_{L^\infty(I, H^r(\Omega))} \\ &+ C\mathcal{M}^{\frac{1}{2\beta\lambda}} h^{\frac{r-1}{2}} \|\rho\|_{L^1(I, W^{r,\beta}(\Omega))}^{\frac{1}{2}} + CT^{\frac{1}{2}} h^r \|\rho_t\|_{L^2(I, H^r(\Omega))}, \end{aligned} \quad (5.2)$$

where $\mathcal{M}$ is defined as in (4.17).

Proof. Subtracting (3.4) from (3.2) we obtain the error equation

$$(\rho_t - \rho_{h,t}, w_h) + (K(|\nabla \rho|)\nabla \rho - K(|\nabla \rho_h|)\nabla \rho_h, \nabla w_h) = 0, \quad w_h \in W_h. \quad (5.3)$$

Choosing $w_h = \theta_h$ in (5.3) gives

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\theta_h\|^2 + (K(|\nabla \rho|)\nabla \rho - K(|\nabla \rho_h|)\nabla \rho_h, \nabla \rho - \nabla \rho_h) \\ = (\vartheta_t, \theta_h) + (K(|\nabla \rho|)\nabla \rho - K(|\nabla \rho_h|)\nabla \rho_h, \nabla \vartheta). \end{aligned} \quad (5.4)$$

By the monotonicity of $K(\cdot)$ in (2.9), there is a positive constant C_0 independence of h and t satisfying

$$(K(|\nabla\rho|)\nabla\rho - K(|\nabla\rho_h|)\nabla\rho_h, \nabla\rho - \nabla\rho_h) \geq C_0\omega(t) \|\nabla\chi\|_{L^\beta}^2.$$

where $\omega(t) = (1 + \|\nabla\rho\|_{L^\beta} + \|\nabla\rho_h\|_{L^\beta})^{-a}$.

Observing from (4.16) that there is a positive constant C_1 independence of h and t such that

$$\begin{aligned} \omega^{-1}(t) &= (1 + \|\nabla\rho\|_{L^\beta} + \|\nabla\rho_h\|_{L^\beta})^{\beta\gamma} \\ &\leq 2^\beta \left(1 + \|\nabla\rho\|_{L^\beta}^\beta + \|\nabla\rho_h\|_{L^\beta}^\beta\right)^\gamma \leq C_1 \mathcal{M}^\gamma. \end{aligned} \quad (5.5)$$

Thus

$$(K(|\nabla\rho|)\nabla\rho - K(|\nabla\rho_h|)\nabla\rho_h, \nabla\rho - \nabla\rho_h) \geq \frac{C_0}{C_1} \mathcal{M}^{-\gamma} \|\nabla\chi\|_{L^\beta}^2. \quad (5.6)$$

By Cauchy's inequality, for $\varepsilon_0 > 0$

$$(\vartheta_t, \theta_h) \leq C\varepsilon_0^{-1} \|\vartheta_t\|^2 + \varepsilon_0 \|\theta_h\|^2, \quad (5.7)$$

Using (2.4), Hölder inequality and then (4.16) we obtain

$$\begin{aligned} (K(|\nabla\rho|)\nabla\rho - K(|\nabla\rho_h|)\nabla\rho_h, \nabla\vartheta) &\leq C(|\nabla\rho|^{\beta-1} + |\nabla\rho_h|^{\beta-1}, |\nabla\vartheta|) \\ &\leq C(\|\nabla\rho\|_{L^\beta}^{\frac{1}{\beta}} + \|\nabla\rho_h\|_{L^\beta}^{\frac{1}{\beta}}) \|\nabla\vartheta\|_{L^\beta} \\ &\leq C(\|\nabla\rho\|_{L^\beta} + \|\nabla\rho_h\|_{L^\beta})^{\frac{1}{\beta}} \|\nabla\vartheta\|_{L^\beta} \\ &\leq C\mathcal{M}^{\frac{1}{\beta\lambda}} \|\nabla\vartheta\|_{L^\beta}. \end{aligned} \quad (5.8)$$

We obtain from (5.4), (5.6), (5.7) and (5.8) that

$$\frac{1}{2} \frac{d}{dt} \|\theta_h\|^2 + \frac{C_0}{C_1} \mathcal{M}^{-\gamma} \|\nabla\chi\|_{L^\beta}^2 \leq C\mathcal{M}^{\frac{1}{\beta\lambda}} \|\nabla\vartheta\|_{L^\beta} + C\varepsilon_0^{-1} \|\vartheta_t\|^2 + \varepsilon_0 \|\theta_h\|^2.$$

Integrating in time from 0 to T and then taking sup-norm in time of resultant show that

$$\|\theta_h\|_{L^\infty(I, L^2)}^2 + \mathcal{M}^{-\gamma} \int_0^T \|\nabla\chi\|_{L^\beta}^2 dt \leq C\mathcal{M}^{\frac{1}{\beta\lambda}} \int_0^T \|\nabla\vartheta\|_{L^\beta} dt + C\varepsilon_0^{-1} \int_0^T \|\vartheta_t\|^2 dt + \varepsilon_0 T \|\theta_h\|_{L^\infty(I, L^2)}^2.$$

Selecting $\varepsilon_0 = \frac{1}{2T}$, we find that

$$\|\theta_h\|_{L^\infty(I, L^2)}^2 + \mathcal{M}^{-\gamma} \int_0^T \|\nabla\chi\|_{L^\beta}^2 dt \leq C\mathcal{M}^{\frac{1}{\beta\lambda}} \int_0^T \|\nabla\vartheta\|_{L^\beta} dt + CT \int_0^T \|\vartheta_t\|^2 dt. \quad (5.9)$$

It follows directly from (5.9) that

$$\begin{aligned} \|\theta_h\|_{L^\infty(I, L^2)} + \mathcal{M}^{-\frac{\gamma}{2}} \|\nabla\chi\|_{L^2(I, L^\beta)} &\leq C\mathcal{M}^{\frac{1}{2\beta\lambda}} \|\nabla\vartheta\|_{L^1(I, L^\beta)}^{\frac{1}{2}} + CT^{\frac{1}{2}} \|\vartheta_t\|_{L^2(I, L^2)} \\ &\leq C\mathcal{M}^{\frac{1}{2\beta\lambda}} h^{\frac{r-1}{2}} \|\rho\|_{L^1(I, W^{r, \beta})}^{\frac{1}{2}} + CT^{\frac{1}{2}} h^r \|\rho_t\|_{L^2(I, H^r)}. \end{aligned} \quad (5.10)$$

By triangle inequality,

$$\|\chi\|_{L^\infty(I, L^2)} \leq \|\vartheta\|_{L^\infty(I, L^2)} + \|\theta_h\|_{L^\infty(I, L^2)}.$$

This and (5.10) imply (5.2). Thus the proof is complete. $\square$

Now we give the error estimate in L^q -norm for any $q > 2$.

THEOREM 5.2. Let $q \in (2, \infty)$, ρ solve the problem (3.2) and ρ_h solve the semidiscrete problem (3.4). Assume that $\rho \in L^\infty(I, W^{r,q}(\Omega) \cap W^{r,2q}(\Omega))$, $\nabla \rho \in L^{2q}(I, L^{2q}(\Omega))$, $\rho_t \in L^q(I, W^{r,q}(\Omega))$. Then there exist constant positive constant C independence of h and T such that

$$\begin{aligned} \|\rho - \rho_h\|_{L^\infty(I, L^q(\Omega))} &\leq \|\rho - \pi\rho\|_{L^\infty(I, L^q(\Omega))} + CT^{1-\frac{1}{q}} \|(\rho - \pi\rho)_t\|_{L^q(I, L^q(\Omega))} \\ &\quad + C\mathcal{A}^{\frac{1}{q}} T^{\frac{1}{2}-\frac{1}{q}} \|\nabla(\rho - \pi\rho)\|_{L^{2q}(I, L^{2q}(\Omega))}, \end{aligned} \quad (5.11)$$

where

$$\mathcal{A} = \left(\int_0^T \left[1 + \|\nabla \rho\|_{L^{aq}(\Omega)}^{aq} + \|\nabla \rho_h\|_{L^{aq}(\Omega)}^{aq} \right] dt \right)^{\frac{1}{2}}. \quad (5.12)$$

Consequently,

$$\begin{aligned} \|\rho - \rho_h\|_{L^\infty(I, L^q(\Omega))} &\leq Ch^r \|\rho\|_{L^\infty(I, W^{r,q}(\Omega))} + CT^{1-\frac{1}{q}} h^r \|\rho_t\|_{L^q(I, W^{r,q}(\Omega))} \\ &\quad + C\mathcal{A}^{\frac{1}{q}} T^{\frac{1}{2}-\frac{1}{q}} h^{r-1} \|\rho\|_{L^{2q}(I, W^{r,2q}(\Omega))}. \end{aligned} \quad (5.13)$$

Proof. Choosing $w_h = |\theta_h|^{q-1} \text{sign} \theta_h$ in (5.3) we have

$$(\vartheta_t - \theta_{h,t}, |\theta_h|^{q-1} \text{sign} \theta_h) + (q-1)(K(|\nabla \rho|) \nabla \rho - K(|\nabla \rho_h|) \nabla \rho_h, |\theta_h|^{q-2} \nabla \theta_h) = 0.$$

We rewrite as form

$$\begin{aligned} &(\theta_{h,t}, |\theta_h|^{q-1} \text{sign} \theta_h) + (q-1)(K(|\nabla \rho|) \nabla \rho - K(|\nabla \rho_h|) \nabla \rho_h, |\theta_h|^{q-2} (\nabla \rho - \nabla \rho_h)) \\ &= (\vartheta_t, |\theta_h|^{q-1} \text{sign} \theta_h) + (q-1)(K(|\nabla \rho|) \nabla \rho - K(|\nabla \rho_h|) \nabla \rho_h, |\theta_h|^{q-2} \nabla \vartheta). \end{aligned}$$

The monotonicity of $K(\cdot)$ in (2.8) provides

$$(K(|\nabla \rho|) \nabla \rho - K(|\nabla \rho_h|) \nabla \rho_h, |\theta_h|^{q-2} (\nabla \rho - \nabla \rho_h)) \geq (\beta-1)(K(\xi) |\nabla \chi|^2, |\theta_h|^{q-2}), \quad (5.14)$$

where $\xi = \max\{|\nabla \rho|, |\nabla \rho_h|\}$. Hence

$$\begin{aligned} &\frac{1}{q} \frac{d}{dt} \|\theta_h\|_{L^q}^q + (q-1)(\beta-1)(K(\xi) |\nabla \chi|^2, |\theta_h|^{q-2}) \\ &\leq (\vartheta_t, |\theta_h|^{q-1} \text{sign} \theta_h) + (q-1)(K(|\nabla \rho|) \nabla \rho - K(|\nabla \rho_h|) \nabla \rho_h, |\theta_h|^{q-2} \nabla \vartheta). \end{aligned}$$

Using Young's inequality with exponents q and $\frac{q}{q-1}$,

$$(\vartheta_t, |\theta_h|^{q-1} \text{sign} \theta_h) \leq q^{-1} \left(\frac{q}{q-1} \right)^{1-q} \varepsilon^{1-q} \|\vartheta_t\|_{L^q}^q + \varepsilon \|\theta_h\|_{L^q}^q. \quad (5.15)$$

Using (2.10), Hölder's inequality and Young's inequality we obtain, for $\varepsilon, \varepsilon_0 > 0$,

$$\begin{aligned} &(K(|\nabla \rho|) \nabla \rho - K(|\nabla \rho_h|) \nabla \rho_h, |\theta_h|^{q-2} \nabla \vartheta) \leq C(|\nabla \chi|, |\theta_h|^{q-2} |\nabla \vartheta|) \\ &\leq \varepsilon_0 (K(\xi) |\nabla \chi|^2, |\theta_h|^{q-2}) + C\varepsilon_0^{-1} (K^{-1}(\xi) |\nabla \vartheta|^2, |\theta_h|^{q-2}) \\ &\leq \varepsilon_0 (K(\xi) |\nabla \chi|^2, |\theta_h|^{q-2}) + \varepsilon \|\theta_h\|_{L^q}^q + C\varepsilon_0^{-\frac{q}{2}} \varepsilon^{-\frac{q-2}{2}} \left\| K^{-\frac{1}{2}}(\xi) \nabla \vartheta \right\|_{L^q}^q. \end{aligned} \quad (5.16)$$

From (5.15), (5.16) with $\varepsilon_0 = (q-1)(\beta-1)/2$, we find that

$$\begin{aligned} &\frac{1}{q} \frac{d}{dt} \|\theta_h\|_{L^q}^q + \frac{(q-1)(\beta-1)}{2} (K(\xi) |\nabla \chi|^2, |\theta_h|^{q-2}) \\ &\leq C\varepsilon^{1-q} \|\vartheta_t\|_{L^q}^q + 2\varepsilon \|\theta_h\|_{L^q}^q + C\varepsilon^{-\frac{q-2}{2}} \left\| K^{-\frac{1}{2}}(\xi) \nabla \vartheta \right\|_{L^q}^q. \end{aligned} \quad (5.17)$$

Integrating in time from 0 to T ,

$$\begin{aligned} & \frac{1}{q} \|\theta_h\|_{L^\infty(I, L^q)}^q + \frac{(q-1)(\beta-1)}{2} \int_0^T (K(\xi) |\nabla \chi|^2, |\theta_h|^{q-2}) dt \\ & \leq 2\varepsilon T \|\theta_h\|_{L^\infty(I, L^q)}^q + C\varepsilon^{1-q} \int_0^T \|\vartheta_t\|_{L^q}^q dt + C\varepsilon^{-\frac{q-2}{2}} \int_0^T \|K^{-\frac{1}{2}}(\xi) \nabla \vartheta\|_{L^q}^q dt. \end{aligned}$$

Choosing $\varepsilon = \frac{1}{4qT}$, using (2.3) we find that

$$\begin{aligned} & \|\theta_h\|_{L^\infty(I, L^q)}^q + \int_0^T (K(\xi) |\nabla \chi|^2, |\theta_h|^{q-2}) ds \\ & \leq CT^{q-1} \int_0^T \|\vartheta_t\|_{L^q}^q dt + CT^{\frac{q-2}{2}} \int_0^T \|K^{-\frac{1}{2}}(\xi) \nabla \vartheta\|_{L^q}^q dt. \end{aligned}$$

Note that

$$\begin{aligned} \int_0^T \|K^{-\frac{1}{2}}(\xi) \nabla \vartheta\|_{L^q}^q dt &= \int_\Omega \int_0^T K^{-\frac{q}{2}}(\xi) |\nabla \vartheta|^q dt dx \\ &\leq \int_\Omega \left(\int_0^T K^{-q}(\xi) dt \right)^{\frac{1}{2}} \left(\int_0^T |\nabla \vartheta|^{2q} dt \right)^{\frac{1}{2}} dx \\ &\leq \left(\int_0^T \int_\Omega K^{-q}(\xi) dx dt \right)^{\frac{1}{2}} \left(\int_0^T \int_\Omega |\nabla \vartheta|^{2q} dx dt \right)^{1/2} \\ &\leq C \left(\int_0^T \int_\Omega (1 + \xi^{aq}) dx dt \right)^{\frac{1}{2}} \|\nabla \vartheta\|_{L^{2q}(I, L^{2q})}^q. \end{aligned}$$

Moreover $1 + \xi^{aq} \leq C(1 + \max\{|\nabla \rho|^{aq}, |\nabla \rho_h|^{aq}\})$. Thus

$$\begin{aligned} \|\theta_h\|_{L^\infty(I, L^q)}^q + \int_0^T (K(\xi) |\nabla \chi|^2, |\theta_h|^{q-2}) dt &\leq CT^{q-1} \int_0^T \|\vartheta_t\|_{L^q}^q dt \\ &\quad + CT^{\frac{q-2}{2}} \left(\int_0^T \int_\Omega 1 + \max\{|\nabla \rho|^{aq}, |\nabla \rho_h|^{aq}\} dx dt \right)^{\frac{1}{2}} \|\nabla \vartheta\|_{L^{2q}(I, L^{2q})}^q. \end{aligned}$$

Dropping the second term of the LHS in above estimate we obtain

$$\|\theta_h\|_{L^\infty(I, L^q)} \leq CT^{1-\frac{1}{q}} \|\vartheta_t\|_{L^q(I, L^q)} + C\mathcal{A}^{\frac{1}{q}} T^{\frac{1}{2}-\frac{1}{q}} \|\nabla \vartheta\|_{L^{2q}(I, L^{2q})}. \quad (5.18)$$

Inequality (5.11) follows thanks to (5.18) and triangle inequality

$$\|\chi\|_{L^\infty(I, L^q)} \leq \|\vartheta\|_{L^\infty(I, L^q)} + \|\theta_h\|_{L^\infty(I, L^q)}.$$

This finishes the proof. $\square$

For the completeness we derive L^∞ -estimate for $\rho - \rho_h$.

THEOREM 5.3. *Let ρ, ρ_h be solutions to problems (3.2) and (3.4) respectively. Assume that $\rho \in L^\infty(I, W^{r,\infty}(\Omega))$, $\rho_t \in L^2(I, W^{r,q}(\Omega))$. Then there exist constant positive constant C independence of h such that*

$$\begin{aligned} \|\rho - \rho_h\|_{L^\infty(I, L^\infty(\Omega))} &\leq Ch^r \|\rho\|_{L^\infty(I, W^{r,\infty}(\Omega))} + CT^{1-\frac{1}{q}} h^{r-\frac{d}{q}} \|\rho_t\|_{L^q(I, W^{r,q}(\Omega))} \\ &\quad + C\mathcal{A}^{\frac{1}{q}} T^{\frac{1}{2}-\frac{1}{q}} h^{r-1-\frac{d}{q}} \|\rho\|_{L^{2q}(I, W^{r,2q}(\Omega))}. \end{aligned} \quad (5.19)$$

Proof. Recall that the quasi-uniform of $\mathcal{T}_h$ we have the inverse estimate (see [4, 35])

$$\|\theta_h\|_{L^\infty(\Omega)} \leq Ch^{-\frac{d}{q}} \|\theta_h\|_{L^q(\Omega)}.$$

This and triangle inequality imply that

$$\begin{aligned} \|\chi\|_{L^\infty(I, L^\infty)} &\leq \|\vartheta\|_{L^\infty(I, L^\infty)} + \|\theta_h\|_{L^\infty(I, L^\infty)} \\ &\leq Ch^r \|\rho\|_{L^\infty(I, W^{r,\infty})} + Ch^{-\frac{d}{q}} \|\theta_h\|_{L^\infty(I, L^q)}. \end{aligned}$$

Thus inequality (5.19) follows directly from (5.18). $\square$

5.2. Error estimate for gradient.

Now we give an error estimate for gradient
THEOREM 5.4. *Under the assumption of Theorem 5.1. For any $0 < t_0 \leq t \leq T$. There exist constant positive constants C independence of h such that*

$$\begin{aligned} \|\nabla(\rho - \rho_h)(t)\|_{L^\beta(\Omega)} &\leq C\mathcal{DM}^{\frac{\gamma}{2}} h^{\frac{r-1}{4}} \left\{ \mathcal{M}^{\frac{1}{4\beta\lambda}} \|\rho(t)\|_{L^2(I, W^{r,\beta}(\Omega))}^{\frac{1}{4}} \right. \\ &\quad \left. + T^{\frac{1}{4}} h^{\frac{r+1}{4}} \|\rho_t(t)\|_{L^2(I, H^r(\Omega))}^{\frac{1}{2}} \right\} + C\mathcal{M}^{\frac{1}{2\beta\lambda}} h^{\frac{r-1}{2}} \|\rho(t)\|_{W^{r,\beta}(\Omega)}^{\frac{1}{2}}. \end{aligned} \quad (5.20)$$

where the positive constant $\mathcal{M}$ is defined as in (4.17), and

$$\mathcal{D} = \left[(t_0^{-1} + 1)\mathcal{M} + \int_0^T h(t) \left(1 + \|f_t(t)\|^2 \right) dt \right]^{\frac{1}{4}}. \quad (5.21)$$

Proof. Choosing $w_h = \theta_h$ in (5.3) we have

$$\begin{aligned} (K(|\nabla\rho|)\nabla\rho - K(|\nabla\rho_h|)\nabla\rho_h, \nabla\rho - \nabla\rho_h) \\ = (\rho_t - \rho_{h,t}, \theta_h) + (K(|\nabla\rho|)\nabla\rho - K(|\nabla\rho_h|)\nabla\rho_h, \nabla\vartheta). \end{aligned} \quad (5.22)$$

By (2.9) and (5.5), there is a positive constant C_0 independence of h such that

$$(K(|\nabla\rho|)\nabla\rho - K(|\nabla\rho_h|)\nabla\rho_h, \nabla\rho - \nabla\rho_h) \geq C_0\mathcal{M}^{-\gamma} \|\nabla(\rho - \rho_h)\|_{L^\beta}^2.$$

In virtue of triangle inequality, Holder's inequality we have

$$\begin{aligned} (\rho_t - \rho_{h,t}, \theta_h) + (K(|\nabla\rho|)\nabla\rho - K(|\nabla\rho_h|)\nabla\rho_h, \nabla\vartheta) \\ \leq (|\rho_t| + |\rho_{h,t}|, |\theta_h|) + C_1(|\nabla\rho|^{\beta-1} + |\nabla\rho_h|^{\beta-1}, |\nabla\vartheta|) \\ \leq (\|\rho_t\| + \|\rho_{h,t}\|) \|\theta_h\| + C_1\mathcal{M}^{\frac{1}{\beta\lambda}} \|\nabla\vartheta\|_{L^\beta} \end{aligned} \quad (5.23)$$

for some positive constant C_1 independence of h .

Thus

$$\|\nabla(\rho - \rho_h)\|_{L^\beta}^2 \leq C_0^{-1}\mathcal{M}^\gamma (\|\rho_t\| + \|\rho_{h,t}\|) \|\theta_h\| + C_1C_0^{-1}\mathcal{M}^{\frac{1}{\beta\lambda}} \|\nabla\vartheta\|_{L^\beta}.$$

Thanks to (4.23) and (4.39),

$$\|\rho_t\| + \|\rho_{h,t}\| \leq 2\mathcal{D}^2. \quad (5.24)$$

Hence there is a positive constant C independence of h such that

$$\|\nabla(\rho - \rho_h)\|_{L^\beta} \leq C\mathcal{DM}^{\frac{\gamma}{2}} \|\theta_h\|^{\frac{1}{2}} + C\mathcal{M}^{\frac{1}{2\beta\lambda}} \|\nabla\vartheta\|_{L^\beta}^{\frac{1}{2}}. \quad (5.25)$$

Due to (5.10) and the fact that $\|\nabla\vartheta\|_{L^\beta} \leq Ch^{r-1} \|\rho\|_{W^{r,\beta}}$ the left hand side of (5.25) is bounded by

$$C\mathcal{DM}^{\frac{\gamma}{2}} \left[\mathcal{M}^{\frac{1}{2\beta\lambda}} h^{\frac{r-1}{2}} \|\rho\|_{L^2(I, W^{r,\beta})}^{\frac{1}{2}} + T^{\frac{1}{2}} h^r \|\rho_t\|_{L^2(I, H^r)} \right]^{\frac{1}{2}} + Ch^{\frac{r-1}{2}} \mathcal{M}^{\frac{1}{2\beta\lambda}} \|\rho\|_{W^{r,\beta}}^{\frac{1}{2}}. \quad (5.26)$$

Thus (5.20) follows from (5.25) and (5.26). $\square$

5.3. Error analysis for fully discrete Galerkin method. In analyzing this method, proceed in a similar fashion as for the semidiscrete method, we derive a error estimate for the fully discrete time Galerkin approximation the differential equation.

Let $\rho^n(\cdot) = \rho(\cdot, t_n)$, be the true solution evaluated at the discrete time levels. We will also denote $\pi\rho^n \in W_h$ to be the projections of the true solutions at the discrete time levels. As in the semidiscrete case, we use $\chi = \rho - \rho_h$, $\vartheta = \rho - \rho_h$, $\theta_h = \rho_h - \pi\rho$ and χ^n , ϑ^n , θ_h^n be evaluating χ , ϑ , θ_h at the discrete time levels. We also define

$$\partial\phi^n = \frac{\phi^n - \phi^{n-1}}{\Delta t}.$$

We rewrite (3.2) with $t = t_n$. Using the definition of L^2 -projection and standard manipulations show that the true solution satisfies the discrete equation

$$(\partial\pi\rho^n, w_h) + (K(|\nabla\rho^n|)\nabla\rho^n, \nabla w_h) = (-\pi\rho_t^n + \partial\pi\rho^n, w_h) - \langle\psi^n, w_h\rangle + (f^n, w_h), \forall w_h \in W_h. \quad (5.27)$$

THEOREM 5.5. *Let ρ^n solve problem (3.2) and ρ_h^n solve the fully discrete Galerkin finite element approximation (3.5) for each time step n , $n = 1, \dots, N$. There exists a positive constant C independent of h and Δt such that if the Δt is sufficiently small then*

$$\max_{1 \leq n \leq N} \|\rho^n - \rho_h^n\| \leq C \left(h^{\frac{r-1}{2}} + \sqrt{\Delta t} \right). \quad (5.28)$$

Proof. Subtracting (5.27) from (3.5) gives the error equation

$$(\partial\theta_h^n, w_h) + (K(|\nabla\rho_h^n|)\nabla\rho_h^n - K(|\nabla\rho^n|)\nabla\rho^n, \nabla w_h) = (\pi\rho_t^n - \partial\pi\rho^n, w_h). \quad (5.29)$$

Selecting $w_h = \theta_h^n$, we rewrite above equation as form

$$\begin{aligned} (\partial\theta_h^n, \theta_h^n) + (K(|\nabla\rho^n|)\nabla\rho^n - K(|\nabla\rho_h^n|)\nabla\rho_h^n, \nabla\rho^n - \nabla\rho_h^n) \\ = (K(|\nabla\rho^n|)\nabla\rho^n - K(|\nabla\rho_h^n|)\nabla\rho_h^n, \nabla\vartheta^n) + (\pi\rho_t^n - \partial\pi\rho^n, \theta_h^n). \end{aligned} \quad (5.30)$$

For the first term, we have the identity

$$\begin{aligned} (\partial\theta_h^n, \theta_h^n) &= \left(\partial\theta_h^n, \frac{\theta_h^n + \theta_h^{n-1}}{2} + \frac{\Delta t}{2} \partial\theta_h^n \right) \\ &= \frac{1}{2\Delta t} \left(\|\theta_h^n\|^2 - \|\theta_h^{n-1}\|^2 \right) + \frac{\Delta t}{2} \|\partial\theta_h^n\|^2. \end{aligned} \quad (5.31)$$

For the second term, the monotonicity of $K(\cdot)$ in (2.9) yields

$$(K(|\nabla\rho^n|)\nabla\rho^n - K(|\nabla\rho_h^n|)\nabla\rho_h^n, \nabla\rho^n - \nabla\rho_h^n) \geq C_0 \omega^n \|\nabla\chi^n\|_{L^\beta}^2 \quad (5.32)$$

with $\omega^n = \omega(t_n) = \left(1 + \max \left\{ \|\nabla\rho_h^n\|_{L^\beta(\Omega)}, \|\nabla\rho^n\|_{L^\beta(\Omega)} \right\} \right)^{-a}$.

For third term, using (2.4), Holder's inequality, there is a positive constant C independence of $h, \Delta t, n$ such that

$$(K(|\nabla\rho^n|)\nabla\rho^n - K(|\nabla\rho_h^n|)\nabla\rho_h^n, \nabla\vartheta^n) \leq C \mathcal{M}^{\frac{1}{\beta\lambda}} \|\nabla\vartheta^n\|_{L^\beta}. \quad (5.33)$$

For the last term, it follows from using L^2 -projection and Taylor expand that

$$(\pi\rho_t^n - \partial\pi\rho^n, \theta_h^n) = (\rho_t^n - \partial\rho^n, \theta_h^n) = \left(\frac{1}{\Delta t} \int_{t_{n-1}}^{t_n} \rho_{tt}(\tau)(\tau - t_{n-1}) d\tau, \theta_h^n \right). \quad (5.34)$$

Differentiating equation (1.15) in time provides,

$$\rho_{tt} = \nabla \cdot (K(|\nabla\rho|)\nabla\rho)_t + f_t.$$

This and (5.34) imply

$$\begin{aligned} (\pi\rho_t^n - \partial\pi\rho^n, \theta_h^n) &= \left(\frac{1}{\Delta t} \int_{t_{n-1}}^{t_n} f_t(\tau)(\tau - t_{n-1}) d\tau, \theta_h^n \right) \\ &\quad - \left(\frac{1}{\Delta t} \int_{t_{n-1}}^{t_n} (K(|\nabla\rho|)\nabla\rho)_t(\tau - t_{n-1}) d\tau, \nabla\theta_h^n \right) = I_1 + I_2. \end{aligned} \quad (5.35)$$

For the first term of left hand side in (5.35),

$$\begin{aligned}
I_1 &\leq \frac{1}{\Delta t} \left\| \int_{t_{n-1}}^{t_n} f_t(\tau)(\tau - t_{n-1}) d\tau \right\| \|\theta_h^n\| \\
&\leq \frac{1}{\Delta t} \|f_t\|_{L^2(I_n, L^2)} \left[\int_{t_{n-1}}^{t_n} (\tau - t_{n-1})^2 d\tau \right]^{\frac{1}{2}} \|\theta_h^n\| \\
&\leq \frac{1}{6} \Delta t \|f_t\|_{L^2(I_n, L^2)}^2 + \frac{1}{2} \|\theta_h^n\|^2,
\end{aligned} \tag{5.36}$$

where $I_n = [t_{n-1}, t_n]$.

For the second term of left hand side in (5.35), using integration by part, triangle inequality, (2.4), and Hölder's inequality we find that

$$\begin{aligned}
I_2 &\leq \frac{1}{\Delta t} \left| \left(\int_{t_{n-1}}^{t_n} K(|\nabla \rho|) \nabla \rho d\tau, \nabla \theta_h^n \right) \right| + |(K(|\nabla \rho^n|) \nabla \rho^n, \nabla \theta_h^n)| \\
&\leq \frac{C}{\Delta t} \int_{t_{n-1}}^{t_n} (|\nabla \rho|^{\beta-1}, |\nabla \theta_h^n|) d\tau + C(|\nabla \rho^n|^{\beta-1}, |\nabla \theta_h^n|) \\
&\leq \frac{C}{\Delta t} \left[\int_{t_{n-1}}^{t_n} \|\nabla \rho\|_{L^\beta}^{\beta-1} d\tau \right] \|\nabla \theta_h^n\|_{L^\beta} + C \|\nabla \rho^n\|_{L^\beta}^{\beta-1} \|\nabla \theta_h^n\|_{L^\beta}.
\end{aligned}$$

According to (4.38),

$$\begin{aligned}
I_2 &\leq C \mathcal{M}^{\frac{\beta-1}{\beta}} \|\nabla \theta_h^n\|_{L^\beta} \\
&\leq C \mathcal{M}^{\frac{\beta-1}{\beta}} \|\nabla \chi^n\|_{L^\beta} + C \mathcal{M}^{\frac{\beta-1}{\beta}} \|\nabla \vartheta^n\|_{L^\beta} \\
&\leq \varepsilon \|\nabla \chi^n\|_{L^\beta}^2 + C \varepsilon^{-1} \mathcal{M}^{2\frac{\beta-1}{\beta}} + C \mathcal{M}^{\frac{\beta-1}{\beta}} \|\nabla \vartheta^n\|_{L^\beta}.
\end{aligned} \tag{5.37}$$

Combining (5.35), (5.36) and (5.37) and using the fact that $\frac{1}{\lambda} = \frac{\beta-1}{\beta}$,

$$\begin{aligned}
(\pi \rho_t^n - \partial \pi \rho^n, \theta_h^n) &\leq \frac{1}{6} \Delta t \|f_t\|_{L^2(I_n, L^2)}^2 + \frac{1}{2} \|\theta_h^n\|^2 \\
&\quad + \varepsilon \|\nabla \chi^n\|_{L^\beta}^2 + C \varepsilon^{-1} \mathcal{M}^{\frac{2}{\lambda}} + C \mathcal{M}^{\frac{1}{\lambda}} \|\nabla \vartheta^n\|_{L^\beta}.
\end{aligned} \tag{5.38}$$

It follows from (5.31), (5.32), (5.33) and (5.38) that

$$\begin{aligned}
\frac{1}{2\Delta t} \left(\|\theta_h^n\|^2 - \|\theta_h^{n-1}\|^2 \right) + C_0 \omega^n \|\nabla \chi^n\|_{L^\beta}^2 &\leq \varepsilon \|\nabla \chi^n\|_{L^\beta}^2 + C(\mathcal{M}^{\frac{1}{\beta\lambda}} + \mathcal{M}^{\frac{1}{\lambda}}) \|\nabla \vartheta^n\|_{L^\beta} \\
&\quad + C \Delta t \|f_t\|_{L^2(I_n, L^2)}^2 + \frac{1}{2} \|\theta_h^n\|^2 + C \varepsilon^{-1} \mathcal{M}^{\frac{2}{\lambda}}.
\end{aligned} \tag{5.39}$$

Selecting $\varepsilon = \frac{C_0 \omega^n}{2}$ in (5.39), we obtain

$$\begin{aligned}
\frac{1}{\Delta t} \left(\|\theta_h^n\|^2 - \|\theta_h^{n-1}\|^2 \right) + C_0 \omega^n \|\nabla \chi^n\|_{L^\beta}^2 &\leq C(\mathcal{M}^{\frac{1}{\beta\lambda}} + \mathcal{M}^{\frac{1}{\lambda}}) \|\nabla \vartheta^n\|_{L^\beta} \\
&\quad + C \Delta t \|f_t\|_{L^2(I_n, L^2)}^2 + \|\theta_h^n\|^2 + C(\omega^n)^{-1} \mathcal{M}^{\frac{2}{\lambda}} + C \mathcal{M}^{\frac{1}{\lambda}} \|\nabla \vartheta^n\|_{L^\beta}.
\end{aligned} \tag{5.40}$$

Dropping the second term in (5.40) and using (5.5) and the fact $\frac{2}{\lambda} + \gamma = 1$, we get

$$\frac{1}{\Delta t} \left(\|\theta_h^n\|^2 - \|\theta_h^{n-1}\|^2 \right) \leq \|\theta_h^n\|^2 + C \left\{ (\mathcal{M}^{\frac{1}{\beta\lambda}} + \mathcal{M}^{\frac{1}{\lambda}}) \|\nabla \vartheta^n\|_{L^\beta} + \mathcal{M}^{\frac{2}{\lambda} + \gamma} \right\} + C \Delta t \|f_t\|_{L^2(I_n, L^2)}^2.$$

Summing from $n = 0$ to $n = m - 1$ with $0 < m \leq N$ and using the fact that $\theta_h(0) = 0$, we obtain

$$\begin{aligned}
\|\theta_h^m\|^2 &\leq \Delta t \sum_{n=0}^{m-1} \|\theta_h^n\|^2 + C \left\{ (\mathcal{M}^{\frac{1}{\beta\lambda}} + \mathcal{M}^{\frac{1}{\lambda}}) \sum_{n=0}^{m-1} \Delta t \|\nabla \vartheta^n\|_{L^\beta} \right. \\
&\quad \left. + (\Delta t)^2 \sum_{n=0}^{m-1} \|f_t\|_{L^2(I_n, L^2)}^2 + (m-1) \mathcal{M} \Delta t \right\}.
\end{aligned} \tag{5.41}$$

An application of the discrete Gronwall's inequality shows that for Δt sufficiently small,

$$\max_{0 \leq n \leq N} \|\theta_h^n\|^2 \leq C \left\{ (\mathcal{M}^{\frac{1}{2\beta\lambda}} + \mathcal{M}^{\frac{1}{\lambda}}) \sum_{n=0}^{m-1} \Delta t \|\nabla \vartheta^n\|_{L^\beta} + \mathcal{M} \Delta t \right\} + C(\Delta t)^2 \|f_t\|_{L^2(I, L^2)}^2, \quad (5.42)$$

which implies

$$\begin{aligned} \max_{0 \leq n \leq N} \|\theta_h^n\| &\leq C \left\{ (\mathcal{M}^{\frac{1}{2\beta\lambda}} + \mathcal{M}^{\frac{1}{\lambda}}) h^{\frac{r-1}{2}} \left(\sum_{n=0}^{N-1} \Delta t \|\rho^n\|_{W^{r,\beta}} \right)^{\frac{1}{2}} + \mathcal{M}^{\frac{1}{2}} \sqrt{\Delta t} \right\} \\ &\quad + C \Delta t \|f_t\|_{L^2(I, L^2)}. \end{aligned} \quad (5.43)$$

Finally, the result follows from (5.43) and $\|\rho^n - \rho_h^n\| \leq \|\theta_h^n\| + \|\vartheta^n\|$. $\square$

Now we give the error estimate for gradient in fully discrete version.

THEOREM 5.6. *Let ρ solve problem (3.2) and ρ_h^n solve the fully discrete Galerkin finite element approximation (3.5) for each time step n , $n = 1, \dots, N$. Then there exists a positive constant C independent of h and Δt satisfying*

$$\|\nabla \rho_h^n - \nabla \rho^n\|_{L^\beta} \leq C(h^{\frac{r-1}{4}} + \sqrt[4]{\Delta t}). \quad (5.44)$$

Proof. Selecting $w_h = \theta_h^n$, in (5.29) and rewriting the resulting equation as form

$$\begin{aligned} ((K(|\nabla \rho^n|)\nabla \rho^n - K(|\nabla \rho_h^n|)\nabla \rho_h^n, \nabla \rho^n - \nabla \rho_h^n) \\ = (K(|\nabla \rho^n|)\nabla \rho^n - K(|\nabla \rho_h^n|)\nabla \rho_h^n, \nabla \vartheta^n) + (\rho_t^n - \partial \rho_h^n, \theta_h^n). \end{aligned}$$

Due to (2.10), Cauchy-Schwartz and triangle inequality, one has

$$\omega^n \|\nabla \rho_h^n - \nabla \rho^n\|_{L^\beta}^2 \leq C (\|\partial \rho_h^n\| + \|\rho_t^n\|) \|\theta_h^n\| + C(|\nabla \rho^n - \nabla \rho_h^n|, |\nabla \vartheta^n|).$$

Since

$$\|\partial \rho_h^n\| = (\Delta t)^{-1} \left\| \int_{t_{n-1}}^{t_n} p_{h,t} dt \right\| \leq (\Delta t)^{-1} \int_{t_{n-1}}^{t_n} \|p_{h,t}\| dt \leq \max_{[T/N, T]} \|p_{h,t}\| \leq \mathcal{D}^2,$$

$\|p_t^n\| \leq \max_{[T/N, T]} \|p_t\| \leq \mathcal{D}^2$ and (5.33), we obtain

$$\omega^n \|\nabla \rho_h^n - \nabla \rho^n\|_{L^\beta}^2 \leq C \mathcal{D}^2 \|\theta_h^n\| + C \mathcal{M}^{\frac{1}{\beta\lambda}} \|\nabla \vartheta^n\|_{L^\beta}.$$

Using estimate in (5.43) and (5.5) shows that

$$\begin{aligned} \|\nabla \rho_h^n - \nabla \rho^n\|_{L^\beta}^2 &\leq C \mathcal{D}^2 \mathcal{M}^\gamma \left\{ (\mathcal{M}^{\frac{1}{2\beta\lambda}} + \mathcal{M}^{\frac{1}{\lambda}}) h^{\frac{r-1}{2}} \left(\sum_{n=0}^{N-1} \Delta t \|\rho^n\|_{W^{r,\beta}} \right)^{\frac{1}{2}} + \mathcal{M}^{\frac{1}{2}} \sqrt{\Delta t} \right. \\ &\quad \left. + \Delta t \|f_t\|_{L^2(I, L^2)} \right\} + C \mathcal{M}^{\gamma + \frac{1}{\beta\lambda}} h^{r-1} \|\rho^n\|_{W^{r,\beta}}, \end{aligned}$$

which proves (5.44). The proof is complete. $\square$

6. Numerical results. In this section, we give a two simple numerical experiments using Galerkin finite element method in the two dimensional region to illustrate the convergent theory. For simplicity, we test the convergence of our method with the Forchheimer two-term law $g(s) = 1 + s$. Equation (1.12) $sg(s) = \xi$, $s \geq 0$ gives $s = \frac{-1 + \sqrt{1 + 4\xi}}{2}$ and hence

$$K(\xi) = \frac{1}{g(s(\xi))} = \frac{2}{1 + \sqrt{1 + 4\xi}}.$$

The region is selected is unit square, i.e $\Omega = [0, 1]^2$. We use the Lagrange element of order $r = 2$ on the unit square in two dimensions. We used FEniCS [29] to perform our numerical

simulations. We divide the unit square into an $N \times N$ mesh of squares, each then subdivide into two right triangles using the `UnitSquareMesh` class in FEniCS. For each mesh, we solve the generalized Forchheimer equation numerically. The error control in each nonlinear solve is $\varepsilon = 10^{-6}$. Our problem is solved at each time level start at $t = 0$ until final time $T = 1$. At this time, we measured the L^2 -errors of pressure and L^β -errors of gradient of pressure and velocity. Here $\beta = 2 - a = 2 - \frac{\deg(g)}{\deg(g)+1} = \frac{3}{2}$.

Example 1. The analytical solution is as follows

$$\rho(x, t) = e^{-2t} \left[\frac{1}{2}(x_1^2 + x_2^2) - \frac{1}{3}(x_1^3 - x_2^3) \right] + 1, \quad \forall (x, t) \in \Omega \times [0, 1].$$

The forcing term f is determined accordingly to the analytical solution by equation $p_t - \nabla \cdot (K(|\nabla \rho|)\rho) = f$. Explicitly,

$$\begin{aligned} f(x, t) = & -2e^{-2t} \left[\frac{1}{2}(x_1^2 + x_2^2) - \frac{1}{3}(x_1^3 - x_2^3) \right] - \frac{4e^{-2t}(1 - x_1 - x_2)}{1 + \sqrt{1 + 4e^{-2t}W(x)}} \\ & + \frac{4e^{-4t} \left[x_1^2(1 - x_1)^2(1 - 2x_1) + x_2^2(1 - x_2)^2(1 - 2x_2) \right]}{W(x) \left(1 + \sqrt{1 + 4e^{-2t}W(x)} \right)^2 \sqrt{1 + 4e^{-2t}W(x)}}. \end{aligned}$$

where $W(x) = \sqrt{x_1^2(1 - x_1)^2 + x_2^2(1 - x_2)^2}$. The initial data $\rho^0(x) = \frac{1}{2}(x_1^2 + x_2^2) - \frac{1}{3}(x_1^3 - x_2^3) + 1$ and the Neumann boundary condition $\psi(x, t) = 0$. The numerical results are listed as the following table.

N	$\ \rho - \rho_h\ $	Rates	$\ \nabla(\rho - \rho_h)\ _{L^\beta(\Omega)}$	Rates
4	6.33E-02	-	4.51E-01	-
8	5.50E-02	0.20	4.07E-01	0.15
16	4.52E-02	0.28	3.70E-01	0.14
32	3.50E-02	0.37	3.22E-01	0.20
64	2.53E-02	0.47	2.70E-01	0.25
128	1.73E-02	0.55	2.21E-01	0.29
256	1.13E-02	0.61	1.79E-01	0.31
512	7.19E-03	0.63	1.43E-01	0.32

Table 1. *Convergence study for generalized Forchheimer flows using Galerkin finite element method with zero flux on the boundary in 2D.*

Example 2. The analytical solution is $\rho(x, t) = x_1 x_2 e^{-t} + 1$ for all $(x, t) \in \Omega \times [0, 1]$. The forcing term f , initial condition and Neumann boundary condition are determined accordingly to the analytical solution as follows

$$f(x, t) = -e^{-t} x_1 x_2 + \frac{8e^{-2t} x_1 x_2}{\sqrt{x_1^2 + x_2^2} \left(1 + \sqrt{1 + 4e^{-t} \sqrt{x_1^2 + x_2^2}} \right)^2 \sqrt{1 + 4e^{-t} \sqrt{x_1^2 + x_2^2}}},$$

and

$$\rho^0(x) = x_1 x_2 + 1, \quad \psi(x, t) = \frac{2e^{-t}}{1 + \sqrt{1 + 4e^{-t} \sqrt{x_1^2 + x_2^2}}} \begin{cases} x_2 & \text{on } x_1 = 0, \\ -x_2 & \text{on } x_1 = 1, \\ -x_1 & \text{on } x_2 = 1, \\ x_1 & \text{on } x_2 = 0. \end{cases}$$

The numerical results are listed in below table

N	$\ \rho - \rho_h\ $	Rates	$\ \nabla(\rho - \rho_h)\ _{L^\beta(\Omega)}$	Rates
4	4.40E-02	-	2.67E-02	-
8	2.24E-02	0.97	2.02E-02	0.40
16	1.15E-02	0.96	1.37E-02	0.56
32	5.90E-03	0.96	8.53E-03	0.68
64	3.01E-03	0.97	4.99E-03	0.77
128	1.53E-03	0.98	2.79E-03	0.84
256	7.70E-04	0.99	1.52E-03	0.88
512	3.94E-4	0.99	8.28E-4	0.92

Table 2. *Convergence study for generalized Forchheimer flow using Galerkin finite element method with nonzero flux on the boundary in 2D.*

Acknowledgment. The author wishes to thank Dr. Luan Hoang for valuable assistance in discussion and suggestions.

REFERENCES

- [1] T. ARBOGAST, M. F. WHEELER, AND N.-Y. ZHANG, *A nonlinear mixed finite element method for a degenerate parabolic equation arising in flow in porous media*, SIAM J. Numer. Anal., 33 (1996), pp. 1669–1687.
- [2] E. AULISA, L. BLOSHANSKAYA, L. HOANG, AND A. IBRAGIMOV, *Analysis of generalized Forchheimer flows of compressible fluids in porous media*, J. Math. Phys., 50 (2009), pp. 103102, 44.
- [3] J. BEAR, *Dynamics of Fluids in Porous Media*, Dover, New York, 1972.
- [4] S. C. BRENNER AND L. R. SCOTT, *The mathematical theory of finite element methods*, vol. 15 of Texts in Applied Mathematics, Springer, New York, third ed., 2008.
- [5] F. BREZZI AND M. FORTIN, *Mixed and hybrid finite element methods*, vol. 15 of Springer Series in Computational Mathematics, Springer-Verlag, New York, 1991.
- [6] P. G. CIARLET, *The finite element method for elliptic problems*, North-Holland Publishing Co., Amsterdam, 1978. Studies in Mathematics and its Applications, Vol. 4.
- [7] E. DiBENEDETTO, *Partial differential equations*, Cornerstones, Birkhäuser Boston Inc., Boston, MA, second ed., 2010.
- [8] J. J. DOUGLAS, P. J. PAES-LEME, AND T. GIORGI, *Generalized Forchheimer flow in porous media*, in Boundary value problems for partial differential equations and applications, vol. 29 of RMA Res. Notes Appl. Math., Masson, Paris, 1993, pp. 99–111.
- [9] K. B. FADIMBA, *Error analysis for a Galerkin finite element method applied to a coupled nonlinear degenerate system of advection-diffusion equations*, Comput. Methods Appl. Math., 6 (2006), pp. 3–30 (electronic).
- [10] ———, *On existence and uniqueness for a coupled system modeling immiscible flow through a porous medium*, J. Math. Anal. Appl., 328 (2007), pp. 1034–1056.
- [11] ———, *A linear backward Euler scheme for the saturation equation: regularity results and consistency*, J. Comput. Appl. Math., 234 (2010), pp. 272–282.
- [12] ———, *A linear backward Euler scheme for a class of degenerate advection-diffusion equations: a mathematical analysis of the convergence in $L^\infty(0, T_0; L^2(\Omega))$ and in $L^2(0, T_0; H^1(\Omega))$* , Anal. Appl. (Singap.), 12 (2014), pp. 227–249.
- [13] K. B. FADIMBA AND R. C. SHARPLEY, *A priori estimates and regularization for a class of porous medium equations*, Nonlinear World, 2 (1995), pp. 13–41.
- [14] ———, *Galerkin finite element method for a class of porous medium equations*, Nonlinear Anal. Real World Appl., 5 (2004), pp. 355–387.
- [15] P. FORCHHEIMER, *Wasserbewegung*, Zeit. Ver. Deut. Ing., 45 (1901), pp. 1781–1788.
- [16] P. FORCHHEIMER, *Hydraulik*, no. Leipzig, Berlin, B. G. Teubner, 1930. 3rd edition.
- [17] L. HOANG AND A. IBRAGIMOV, *Structural stability of generalized Forchheimer equations for compressible fluids in porous media*, Nonlinearity, 24 (2011), pp. 1–41.
- [18] L. HOANG AND A. IBRAGIMOV, *Qualitative Study of Generalized Forchheimer Flows with the Flux Boundary Condition*, Adv. Diff. Eq., 17 (2012), pp. 511–556.
- [19] L. HOANG, A. IBRAGIMOV, T. KIEU, AND Z. SOBOLE, *Stability of solutions to generalized Forchheimer equations of any degree*, J. Math. Sci., (2015). accepted.
- [20] L. HOANG AND T. KIEU, *Global estimates for generalized Forchheimer flows of slightly compressible fluids*, Journal d'Analyse Mathématique, (2015). accepted.
- [21] ———, *Interior estimates for generalized Forchheimer flows of slightly compressible fluids*, (2015). submitted, preprint <http://arxiv.org/abs/1404.6517>.
- [22] L. T. HOANG, A. IBRAGIMOV, AND T. T. KIEU, *One-dimensional two-phase generalized Forchheimer flows of incompressible fluids*, J. Math. Anal. Appl., 401 (2013), pp. 921–938.
- [23] ———, *A family of steady two-phase generalized Forchheimer flows and their linear stability analysis*, J. Math. Phys., 55 (2014), p. 123101.

- [24] L. T. HOANG, T. T. KIEU, AND T. V. PHAN, *Properties of generalized Forchheimer flows in porous media*, J. Math. Sci., 202 (2014), pp. 259–332.
- [25] C. JOHNSON AND V. THOMÉE, *Error estimates for some mixed finite element methods for parabolic type problems*, RAIRO Anal. Numér., 15 (1981), pp. 41–78.
- [26] T. KIEU, *Analysis of expanded mixed finite element methods for the generalized Forchheimer flows of slightly compressible fluids*, Numer. Methods Partial Differential Equations, (2015). accepted.
- [27] O. A. LADYŽENSKAJA, V. A. SOLONNIKOV, AND N. N. URALCÉVA, *Linear and quasilinear equations of parabolic type*, Translated from the Russian by S. Smith. Translations of Mathematical Monographs, Vol. 23, American Mathematical Society, Providence, R.I., 1968.
- [28] J.-L. LIONS, *Quelques méthodes de résolution des problèmes aux limites non linéaires*, Dunod, 1969.
- [29] A. LOGG, K.-A. MARDAL, AND G. N. WELLS, eds., *Automated Solution of Differential Equations by the Finite Element Method*, vol. 84 of Lecture Notes in Computational Science and Engineering, Springer, 2012.
- [30] M. MUSKAT, *The flow of homogeneous fluids through porous media*, McGraw-Hill Book Company, inc., 1937.
- [31] D. A. NIELD AND A. BEJAN, *Convection in porous media*, Springer-Verlag, New York, second ed., 1999.
- [32] E.-J. PARK, *Mixed finite element methods for generalized Forchheimer flow in porous media*, Numer. Methods Partial Differential Equations, 21 (2005), pp. 213–228.
- [33] R. E. SHOWALTER, *Monotone operators in Banach space and nonlinear partial differential equations*, vol. 49 of Mathematical Surveys and Monographs, American Mathematical Society, Providence, RI, 1997.
- [34] B. STRAUGHAN, *Stability and wave motion in porous media*, vol. 165 of Applied Mathematical Sciences, Springer, New York, 2008.
- [35] V. THOMÉE, *Galerkin finite element methods for parabolic problems*, vol. 25 of Springer Series in Computational Mathematics, Springer-Verlag, Berlin, second ed., 2006.
- [36] J. C. WARD, *Turbulent flow in porous media.*, Journal of the Hydraulics Division, Proc. Am. Soc. Civ. Eng., 90(HY5) (1964), pp. 1–12.
- [37] C. S. WOODWARD AND C. N. DAWSON, *Analysis of expanded mixed finite element methods for a nonlinear parabolic equation modeling flow into variably saturated porous media*, SIAM J. Numer. Anal., 37 (2000), pp. 701–724 (electronic).
- [38] E. ZEIDLER, *Nonlinear functional analysis and its applications. II/B*, Springer-Verlag, New York, 1990. Nonlinear monotone operators, Translated from the German by the author and Leo F. Boron.